

2017-06

# The challenges and opportunities of addressing particle size effects in sediment source fingerprinting: A review

Laceby, JP

<http://hdl.handle.net/10026.1/9622>

---

10.1016/j.earscirev.2017.04.009

Earth-Science Reviews

Elsevier BV

---

*All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.*

# The challenges and opportunities of addressing particle size effects in sediment source fingerprinting: a review

J. Patrick Lacey<sup>a\*</sup>, Olivier Evrard<sup>a</sup>, Hugh G. Smith<sup>b</sup>, Will H. Blake<sup>c</sup>, Jon M. Olley<sup>d</sup>, Jean P. G. Minella<sup>e</sup>  
Philip N. Owens<sup>f</sup>

<sup>a</sup> Laboratoire des Sciences du Climat et de l'Environnement (LSCE/IPSL), Unité Mixte de Recherche 8212 (CEA/CNRS/UVSQ), Université Paris-Saclay, Gif-sur-Yvette, 91198, France

<sup>b</sup> School of Environmental Sciences, University of Liverpool, Liverpool, L69 7ZT, UK

<sup>c</sup> School of Geography, Earth and Environmental Sciences, Plymouth University, Plymouth, PL4 8AA, UK

<sup>d</sup> Australian Rivers Institute, Griffith University, 170 Kessels Road, Nathan, QLD 4101, Australia

<sup>e</sup> Universidade Federal de Santa Maria, Department of Soils, 1000 Roraima Avenue, 97105-900 Santa Maria, RS, Brazil

<sup>f</sup> Environmental Science Program and Quesnel River Research Centre, University of Northern British Columbia, Prince George, British Columbia, V2N 4Z9, Canada

## Abstract:

Tracing sediments back to their catchment sources using biogeochemical and physical fingerprints involves multiple assumptions. One of the most fundamental assumptions is that these fingerprints are consistent during sediment generation, transportation, and deposition processes. Accordingly, the biogeochemical fingerprints used to trace sediment must remain constant, during detachment and redistribution, or they must vary in a predictable and measurable way. One key challenge to this assumption is the sorting effect of particles by size during detachment, mobilization, transportation and deposition processes. Owing to the notable effect of particle size on sediment fingerprints, we believe it is important to review the main approaches used to address the effects of changes in particle size composition on sediment fingerprints. The two main approaches to addressing particle size impacts on fingerprint properties are: fractionation of source and sediment material to a narrow particle size range (e.g. isolation of <10 µm or <63 µm fractions), and concentration corrections (e.g. normalising concentrations by parameters such as specific surface area). These approaches are often used in combination. The utility of fractionation and corrections to address particle size effects has received increasing attention and the relative merits of these procedures have been subject to debate. Accordingly, alternative techniques to address particle size effects in sediment fingerprinting studies are being adopted. For example, a tributary tracing technique or edge-of-field samplers may minimise particle size effects on sediment source fingerprints. The interrelationships between particle size and biogeochemical tracer properties suggest that particle size may also contribute to the formation of contrasts in sediment fingerprints between sources. Indeed, there may be a significant opportunity to derive further sediment source information through comprehensively investigating and unravelling the complexity of particle size–biogeochemical interactions.

**Key words:** Grain size; sediment fingerprinting; composite fingerprinting; sediment tracing; sediment provenance

## \*Corresponding author:

J. Patrick Lacey

Phone: +33 07 82 35 35 48

Fax: +33 01 69 82 35 68

E-mail: [placeby@lsce.ipsl.fr](mailto:placeby@lsce.ipsl.fr)

**Article in *Earth-Science Reviews* 169 · Accepted 18th April 2017**

**DOI: 10.1016/j.earscirev.2017.04.009**

**Embargo: 24 months**

## 1. Introduction

Accelerated soil erosion impacts land and water quality worldwide. Although sediment is a natural component of fluvial systems that provides fundamental structure to riverine landscapes and is essential in many aquatic ecosystems (Vercruysse et al., 2017; Wohl et al., 2015), the excess supply of fine sediment from accelerated soil erosion often degrades riverine and coastal environments (McCulloch et al., 2003; Owens et al., 2005; Walling and Collins, 2016) and contributes to the downstream transfer of particle-bound contaminants (Gateuille et al., 2014; Yamashiki et al., 2014). Elevated suspended sediment loads may also increase the cost of operating and maintaining water treatment and transportation infrastructure (Clark, 1985). Knowledge of the relative contribution of different sources supplying sediment to riverine, lacustrine and coastal systems is a crucial prerequisite to implementing efficient best practices necessary to limit the off-site impacts of excessive sediment delivery (Belmont et al., 2011; Koiter et al., 2013b).

One increasingly adopted field-based approach to identifying sources supplying material to riverine, lacustrine and coastal environments is sediment fingerprinting. Tracing sediments back to their primary sources with fingerprinting techniques offers a direct method to identify the nature, location and relative source contribution of sediment transported in waterways. Sediment properties such as mineral magnetic parameters, fallout radionuclides, major and trace element geochemistry, and compound specific stable isotopes (CSSI) have all been used to trace sediment sources (Blake et al., 2012; Caitcheon, 1993; Evrard et al., 2011; Hancock and Revill, 2013; Murray et al., 1993b; Walling and Kane, 1984).

For properties to be effective tracers of sediment, they must differentiate between sediment sources whilst behaving conservatively (Walling et al., 1993). Conservative behavior is characterized by constancy in sediment properties, where the properties of sediment sources remain constant, or at the very least, any variation in these properties should occur in a predictable and measurable way. In addition, properties of the eroded sediment should remain constant through sediment detachment, transportation and deposition processes, or again, vary in a predictable and measurable way (Belmont et al., 2014; Koiter et al., 2013b; Motha et al., 2002; Olley et al., 2001).

Erosion and transport processes are selective regarding the particle size of the material affected. Detachment processes that generate sediment for fluvial transport are particle size dependent. Clay particles may resist detachment depending on the strength of their bond with the substrate whereas coarse sand may resist detachment simply as a result of size and weight (Bradford et al., 1992; Poesen,

1992). Silt and fine sand are thus more subject to detachment and subsequent transport as they are lighter and without bonds binding them to the substrate (Morgan, 2005; Poesen, 1992). Thereafter, the particle size of suspended sediment directly influences settling velocities in aquatic systems (Gibbs et al., 1971) resulting in the transport and deposition of suspended sediment being particle size selective (Viparelli et al., 2013; Walling et al., 2000). Fluvial transport, including overland flow, produces changes in the characteristics of the material being transported in comparison to the original source material. In general, the average size of particles decreases, while the degree of sorting and the average roundness increases, with distance travelled. These changes result from a combination of selective transportation, deposition, and particle abrasion, with these processes acting over the entire landscape (Frings, 2008; Krumbein and Sloss, 1951; Le et al., 2015; Moss and Walker, 1978).

Not only are sediment transport processes particle-size selective, the properties used to trace sediments may have different affinities to various particle size fractions. For example, fallout radionuclides are preferentially bound to clay minerals owing to the higher number of potential sorption sites (Fan et al., 2014; Lomenick and Tamura, 1965; Tamura, 1964). Magnetic minerals occur in soil and sediment as aggregated concretions, discrete fine grains, and particle coatings on very fine grains (Oldfield, 1991; Smith, 1999). Different geochemical elements are contained within the mineral matrix or adsorbed (Stumm and Morgan, 2012). Organic matter may coat grains or be bound within the mineral matrix (Keil and Mayer, 2014; Mayer, 1999). Ultimately, the properties used to trace sediment may have different affinities (e.g. preferential adsorption/absorption) for various particles size fractions that in combination with the selective transport of fine-grained material may affect sediment source fingerprinting results.

While there are multiple literature reviews published on sediment source fingerprinting (Collins and Walling, 2002; D'Haen et al., 2012; Davis and Fox, 2009; Guzmán et al., 2013; Haddadchi et al., 2013; Koiter et al., 2013b; Owens et al., 2016; Smith et al., 2013), and a recent emphasis on modelling approaches (Cooper et al., 2014; Haddadchi et al., 2014; Laceby and Olley, 2015; Palazón et al., 2015b; Zhang and Liu, 2016), few studies have investigated the effects of particle size on sediment source signatures (Olley and Murray, 1994; Russell et al., 2001; Smith and Blake, 2014). Here, we review the effects and challenges (section 2), approaches (sections 3 and 4) and opportunities (section 5) of particle size selectivity within the sediment source fingerprinting technique. We mainly focus on riverine environments, specifically particle size selectivity as sediment moves from hillslopes into and through river channels, but the concepts and examples presented are also relevant for similar applications in other aquatic systems such as lakes, reservoirs, estuaries and the coastal zone.

## 2. Sediment Property Predictability

Sediment generation, transportation and deposition processes are known to be particle size selective, where fine particles generally have a greater probability of being detached and transported further than coarse particles (McLaren and Bowles, 1985; Walling and Moorehead, 1989). This particle size selectivity often results in potential differences in biogeochemical tracer property concentrations in detached material relative to their sources (He and Walling, 1996; Horowitz and Elrick, 1987). Accordingly, this section will review the impacts of particle size selectivity on a variety of biogeochemical properties used in sediment source fingerprinting research. We focus on some of the main properties – such as fallout radionuclides, carbon and nitrogen parameters, elemental geochemistry and mineral magnetic properties – but recognise that many of the findings are also likely to be relevant to other biogeochemical properties used as fingerprints such as colour parameters and DNA.

### 2.1 Fallout Radionuclides ( $^{137}\text{Cs}$ , $^{210}\text{Pb}_{\text{ex}}$ , $^7\text{Be}$ )

Caesium-137 ( $^{137}\text{Cs}$ ,  $T_{1/2}= 30$  y) and excess lead-210 ( $^{210}\text{Pb}_{\text{ex}}$ ,  $T_{1/2}= 22$  y) have been widely used to determine the relative contributions of sediment from different erosion processes to waterways (Ben Slimane et al., 2016; IAEA, 2014; Matisoff et al., 2002; Owens et al., 2012; Smith et al., 2011; Wallbrink et al., 1998; Walling and Woodward, 1992) (Table 1). As both  $^{137}\text{Cs}$  and  $^{210}\text{Pb}_{\text{ex}}$  are concentrated near the soil surface, as they are atmospheric fallout products, sediments eroded from rill or sheet erosion often have high  $^{137}\text{Cs}$  and  $^{210}\text{Pb}_{\text{ex}}$  activity concentrations (Walling, 2005), whereas sediments eroded from subsoil channel bank or gully erosion processes have low  $^{137}\text{Cs}$  and  $^{210}\text{Pb}_{\text{ex}}$  activity concentrations (Belmont et al., 2014; Olley et al., 2013; Wallbrink et al., 1999). Comparing  $^{137}\text{Cs}$  and  $^{210}\text{Pb}_{\text{ex}}$  in suspended sediments and sediments generated by these different erosion processes generally allows for the relative sediment contributions from these different erosion processes to be ascertained (Ben Slimane et al., 2013; Wallbrink and Murray, 1993; Wallbrink et al., 1999; Walling, 2003).

**Table 1:** Examples of research utilizing only fallout radionuclides to trace sediment sources (SSA refers to specific surface area).

Reference	Country	Area (km <sup>2</sup> )	Parameters	Particle Size (µm)	Corrections or other approaches to address particle size dynamics	Pre-treatment before analysis	Raw Data Availability
Belmont et al., 2011	United States	2880	<sup>10</sup> Be, <sup>210</sup> Pb <sub>ex</sub> , <sup>137</sup> Cs	<64, <125 <sup>a</sup>	SSA	--	Yes
Belmont et al., 2014	United States	880	<sup>10</sup> Be, <sup>210</sup> Pb <sub>ex</sub> , <sup>137</sup> Cs	<64, <125 <sup>a</sup>	--	--	Yes
Blake et al., 2009	Australia	<1	<sup>7</sup> Be, <sup>210</sup> Pb <sub>ex</sub> , <sup>137</sup> Cs	8 fractions <sup>b</sup>	Enrichment factors	Dispersed <sup>c</sup>	Partial <sup>d</sup>
Bonniwell et al., 1999	United States	389	<sup>7</sup> Be, <sup>210</sup> Pb <sub>ex</sub> , <sup>137</sup> Cs	--	Ratios	--	Yes
Evrard et al., 2010	Mexico	3, 9, 12	<sup>7</sup> Be, <sup>210</sup> Pb <sub>ex</sub> , <sup>137</sup> Cs	--	Sc Comparison	--	Partial <sup>e</sup>
Evrard et al., 2016	Laos	12	<sup>7</sup> Be, <sup>210</sup> Pb <sub>ex</sub> , <sup>137</sup> Cs	<1000	Runoff samplers	--	Yes
Foucher et al., 2015	France	24	<sup>137</sup> Cs	<20, 20-50, 50-63, <2000	Th correction, SSA	--	Partial <sup>e</sup>
He and Owens, 1995	United Kingdom		<sup>226</sup> Ra, <sup>210</sup> Pb <sub>ex</sub> , <sup>137</sup> Cs	<2000	SSA	--	Yes
Gourdin et al., 2014	Laos	12	<sup>7</sup> Be, <sup>210</sup> Pb <sub>ex</sub> , <sup>137</sup> Cs	<1000	Ratios	--	Partial <sup>e</sup>
Matisoff et al., 2002	United States	70	<sup>7</sup> Be, <sup>210</sup> Pb <sub>ex</sub> , <sup>137</sup> Cs	--	--	--	--
Matisoff et al., 2005	United States	--	<sup>7</sup> Be, <sup>210</sup> Pb <sub>ex</sub>	--	Ratios	--	--
Murray et al., 1993a	Australia	--	<sup>137</sup> Cs, <sup>226</sup> Ra, <sup>232</sup> Th	<2000	--	--	--
Olley et al., 2013	Australia	47-3842	<sup>210</sup> Pb <sub>ex</sub> , <sup>137</sup> Cs	<10	--	--	--
Owens et al., 2012	Canada	135, 215	<sup>210</sup> Pb <sub>ex</sub> , <sup>137</sup> Cs	<63	SSA	--	--
Smith et al., 2011	Australia	1.4	<sup>210</sup> Pb <sub>ex</sub> , <sup>137</sup> Cs	<63	SSA	--	Yes
Stout et al., 2014	United States	4,300	<sup>10</sup> Be, <sup>210</sup> Pb <sub>ex</sub> , <sup>137</sup> Cs	<64, <125 <sup>a</sup>	SSA	--	--
Wallbrink and Murray, 1993	Australia	<0.001	<sup>7</sup> Be, <sup>210</sup> Pb <sub>ex</sub> , <sup>137</sup> Cs	--	--	--	--
Wallbrink et al., 1998	Australia	13500	<sup>210</sup> Pb <sub>ex</sub> , <sup>137</sup> Cs	<2 <sup>a</sup>	--	--	--
Walling and Woodward, 1992	United Kingdom	12, 46	<sup>7</sup> Be, <sup>210</sup> Pb <sub>ex</sub> , <sup>137</sup> Cs	<63	Source correction	--	--

<sup>a</sup> Source samples were sieved to a size fraction that matched sediment samples

<sup>b</sup> Fractions: <10, 10-20, 20-40, 40-63, 63-125, 125-250, 250-500, 500-2000µm

<sup>c</sup> Sonified before sieving prior to analyses

<sup>d</sup> For outwash sediment sample


<sup>e</sup> Sediments not soils

Although they have been used for tracing sediment generated from erosion processes (Wallbrink and Murray, 1996b), beryllium-7 ( $^7\text{Be}$ ,  $T_{1/2} = 53$  d) and  $^{210}\text{Pb}_{\text{ex}}$  are also increasingly used as chronometers of sediment transfers in riverine systems (Bonniwell et al., 1999; Evrard et al., 2016; Gourdin et al., 2014; Mabit et al., 2014; Smith et al., 2014; Taylor et al., 2013). For example, these radionuclides quantify the relative sediment contribution from 'old' ( $^7\text{Be}$ -depleted) and 'new' ( $^7\text{Be}$ -enriched) sources (Evrard et al., 2010; Matisoff et al., 2005). Over longer time scales, additional tracers are capable of providing further chronological information (e.g.  $^{10}\text{Be}$ ,  $T_{1/2} = 1.39 \times 10^6$  y) (Belmont et al., 2011; Stout et al., 2014). In particular, Belmont et al. (2014) combined  $^{10}\text{Be}$ ,  $^{210}\text{Pb}_{\text{ex}}$  and  $^{137}\text{Cs}$  measurements to demonstrate the potential for over-estimating channel source contributions when there is a moderate amount of sediment exchange between the channel and the floodplain in large watersheds over sediment routing timescales.

Research has demonstrated that fallout radionuclides are typically enriched in the fine particle size fractions. For example, He and Walling (1996) reported increasing activity concentrations of  $^{137}\text{Cs}$  and  $^{210}\text{Pb}_{\text{ex}}$  with increasing specific surface area (SSA) (Figure 1). SSA is closely related to particle size (Horowitz, 1991) and is reported as the total surface area per unit mass (Rawlins et al., 2010). For example, the SSA of sediment increases with decreasing particle size to the extent that the SSA values for clays may be several orders of magnitude greater than silt and sand (Walling and Moorehead, 1989). As fallout radionuclides are preferentially bound to clay minerals owing to the higher number of potential sorption sites (Fan et al., 2014; Lomenick and Tamura, 1965; Tamura, 1964), activity concentrations typically increase with increasing SSA and decreasing particle size (He and Owens, 1995; Wallbrink et al., 1999). Although less research has documented the relationship between particle size and  $^7\text{Be}$ , this fallout radionuclide has also been found to be enriched in fine particle size fractions (Blake et al., 2009; Taylor et al., 2014; Wallbrink and Murray, 1996a). As fallout radionuclides are typically enriched in fine particle size fractions, it may be possible to quantify their relationship with SSA.

The challenge is that there are exceptions to the rule. For example, Smith and Blake (2014) observed that  $^{137}\text{Cs}$  and  $^{210}\text{Pb}_{\text{ex}}$  were negatively related to SSA in pasture soils ( $p < 0.05$  for  $^{210}\text{Pb}_{\text{ex}}$  only), whereas they exhibited positive though non-significant relationships with SSA in channel bank and cultivated soils. This highlights the fact that although these fallout radionuclides are generally enriched in the fine particle size fractions, there may be exceptions, and individual sources may behave differently. Potential exceptions may be explained by the presence and amount of HCl-extractable materials, which Singleton et al. (2017) reported to have a stronger control on fallout radionuclides than grain size or mineralogy. Furthermore, the depth-dependent distribution of fallout radionuclides in the soil profile relates to the exposure to

fallout and subsequent diffusion and migration processes (Jagercikova et al., 2015). Therefore, fallout radionuclide activity concentrations may decrease with soil depth despite increasing clay content as deeper soil was not exposed to fallout. These exceptions demonstrate the need to understand the relationship between particle size distribution and the tracer property of interest in each study.



**Figure 1:** Relationship between specific surface area and fallout radionuclide activity concentrations (<sup>137</sup>Cs and <sup>210</sup>Pb<sub>ex</sub>) (adaptation of Figure 1 from He and Walling (1996)).

## 2.2 Carbon and Nitrogen Parameters

Although they are not as extensively analyzed in sediment tracing research as fallout radionuclides, carbon and nitrogen parameters provide an interesting example into tracer property relationships with particle size. Total organic carbon (TOC) and total nitrogen (TN) often discriminate between sediment derived from surface and subsoil erosion processes as their concentration decreases with depth in the soil profile (Blake et al., 2006; Owens et al., 2006). Carbon stable isotopes ( $\delta^{13}\text{C}$ ) may potentially discriminate between sediment derived from soils with C<sub>3</sub> vegetation (majority of tree or temperate grass species) compared to those covered with C<sub>4</sub> vegetation (grass and cropping species typically under warmer climates) (Fry, 2006; Schimel, 1993). Source discrimination with nitrogen stable isotopes ( $\delta^{15}\text{N}$ ) is more complex. In general, soil  $\delta^{15}\text{N}$  increases with depth in soil profiles (Amundson et al., 2003; Natelhoffer and Fry, 1988). Other factors such as nitrogen inputs from animal and human waste, along with fertilizers and potentially topographic position, may impact the predictability of  $\delta^{15}\text{N}$  in a tracing context. Of note, these parameters trace organic matter (Garzon-Garcia et al., 2017; Olley, 2002) and therefore they may be fractionated by density as well as particle size. Furthermore, these parameters also have the potential for non-conservative behavior due to biological uptake and consumption. Nonetheless, these parameters are increasingly used in sediment



tracing research (Fox and Papanicolaou, 2007; Laceby et al., 2015b; Mukundan et al., 2010; Papanicolaou et al., 2003) (Table 2).

Similarly to fallout radionuclides, TOC and TN are generally enriched in the fine particle size fraction (Balesdent et al., 1987; Wynn et al., 2005). For example, when normalizing the <2, <10, and <63  $\mu\text{m}$  fractions by the bulk soil (<2 mm) fraction, Laceby et al. (2015b) reported that the <63  $\mu\text{m}$  fraction was significantly different than the <2 and <10  $\mu\text{m}$  fractions for TN, though not for TOC, indicating that these similar properties may behave slightly differently in two Australian catchments (Figure 2). Laceby et al. (2016) also reported significant enrichment for TOC and TN between the bulk soil and the <63  $\mu\text{m}$  fraction for subsoils and cultivated sources, though not for forest source samples in several Japanese catchments (Figure 3).


More variability is anticipated in the relationship between particle size and  $\delta^{15}\text{N}$  and  $\delta^{13}\text{C}$  (Balesdent et al., 1987; Bellanger et al., 2004). However, Laceby et al. (2015b) reported that there were not significant differences between particle size fractions for  $\delta^{15}\text{N}$ , with very limited  $\delta^{13}\text{C}$  variation across different particle size fractions (Figure 2). Similarly, Laceby et al. (2016) found little variation between the bulk soil fraction and the <63  $\mu\text{m}$  fraction for  $\delta^{13}\text{C}$ , with increasing, though not significant, enrichment for  $\delta^{15}\text{N}$  (Figure 3).

Based on the fundamental principles of sediment source fingerprinting, as long as the enrichment is predictable, these organic sediment properties could potentially be effective tracers of erosion processes, particularly in contexts where fallout radionuclides are ineffective.  $\delta^{15}\text{N}$  appears more complex and may be enriched or depleted in the different particle size fractions, which requires investigation on a case by case basis.  $\delta^{13}\text{C}$  apparently behaves very predictably in a sediment tracing context although there will likely be exceptions. Importantly, the particle size enrichment for TOC, TN and  $\delta^{15}\text{N}$  varied for the individual sources (Laceby et al., 2016). Therefore, it is crucial to understand the predictability of particle size effects when tracing sediment sources with these and other carbon and nitrogen properties (e.g. compound specific stable isotopes (Reiffarth et al., 2016)).


**Table 2:** Examples of research using only carbon (C) and nitrogen (N) parameters to trace sediment sources

Reference	Country	Area (km <sup>2</sup> )	Parameters	Particle Size (µm)	Corrections or other approaches to address particle size dynamics	Pre-treatment before analysis	Raw Data Available
Fox and Papanicolaou, 2007	United States	0.71	$\delta^{13}\text{C}$ , $\delta^{15}\text{N}$ , C/N	<53	--	Dispersed	Yes
Garzon-Garcia et al., 2017	Australia	2.5, 75, 3076	$\delta^{13}\text{C}$ , $\delta^{15}\text{N}$ , TOC, TN	<10, <63, <500	--	--	Yes
Gibbs, 2008	New Zealand	117	CSSI	<1000	--	--	Yes
Hancock and Revill, 2013	Australia	3860	CSSI	<63	--	--	--
Lacey et al., 2015b	Australia	75, 123, 311	$\delta^{13}\text{C}$ , $\delta^{15}\text{N}$ , TOC, TN	<2, <10, <63, <2000	--	--	--
Lacey et al., 2016	Japan	77, 171, 265	$\delta^{13}\text{C}$ , $\delta^{15}\text{N}$ , TOC, TN	<63, <2000	--	--	Yes
Olley, 2002	Australia	84000	$\delta^{13}\text{C}$ , C/N Ratio	<2	--	--	--
Papanicolaou et al., 2003	United States	600	$\delta^{13}\text{C}$ , $\delta^{15}\text{N}$ , C/N	--	--	--	--

<sup>a</sup> Source samples were dispersed with sodium hexametaphosphate



**Figure 2:** Normalized carbon (C) and nitrogen (N) property distributions for the <2  $\mu\text{m}$ , <10  $\mu\text{m}$  and <63  $\mu\text{m}$  particle size fractions in samples from Knapp Creek and Blackfellow Creek, Australia (modified from Laceby et al., (2015b)). Points under the distributions are the normalized samples color coded to particle size fraction used to derive these distributions with kernel density functions.



**Figure 3:** Box plots of the normalized difference between carbon (C) and nitrogen (N) sediment properties in the  $<63\ \mu\text{m}$  fraction divided by the  $<2\ \text{mm}$  fraction for the three sources and sediment from three Japanese catchments (i.e., Mano, Niida and Ota) with red shading (and the red circle) indicating significant differences between the two fractions and values greater than 1 (the dotted line) being enriched in the  $<63\ \mu\text{m}$  fraction (modified from Laceby et al., (2016)).

## 2.3 Elemental Geochemistry

Major and trace elemental geochemistry (including rare earth elements) are often used to identify the different spatial sources of sediment (Hardy et al., 2010; Laceby and Olley, 2015; Vale et al., 2016) (Table 3). Different parent rock material typically results in sources having distinct elemental geochemistry (Douglas et al., 2009; Motha et al., 2002; Olley et al., 2001). Eroded sediment often maintain these geochemical fingerprints, allowing the relative contributions of different sources to be ascertained (Caitcheon et al., 2006; D'Haen et al., 2013; Hughes et al., 2009). The question is whether these fingerprints are conservative during sediment generation, transportation and deposition processes.

The relationship between particle size enrichment and elemental geochemistry is complex and partly dependent on the digestion procedure (e.g. acid leached versus total digestion) used to prepare samples

for analysis (e.g. inductively coupled plasma mass spectrometry (ICP-MS)). The difficulty is that elemental analyses often provide results for over 40 elements (Table 3) whereas the previous sections examined three fallout radionuclides and four carbon and nitrogen parameters. For each of these 40 plus elements, sediment generation, transport and deposition processes may potentially enrich their elemental concentrations, deplete them, or have a limited impact (e.g. Motha et al., 2002; Russell et al., 2001; Smith and Blake, 2014). The impact of particle size selectivity on elemental geochemistry likely will depend on how elements are incorporated into fine sediment (e.g. within the mineral matrix or adsorbed).

Underlying the potential influence of particle size on elemental geochemistry is the effect of sediment source mineralogy. The dominant mineralogy relates directly to particle size where some sources will be enriched in the fine particle size fractions and other sources may be depleted. Thus the geochemical fingerprint likely will change if the <2 mm, the <63  $\mu\text{m}$ , or the <10  $\mu\text{m}$  fraction are sampled and analysed. For example, each of the three different sources in Figure 4 from Gibbs (1967) may have different relationships between elemental concentrations and particle size, which will depend on the particle size range utilized for the sediment source fingerprinting research. Fundamentally, the impact of particle size on all elemental concentrations for each source is difficult to predict and the complexity of the particle size – elemental geochemistry relationship requires more research to comprehensively characterize the predictability of these fingerprints. Accordingly, section 4 of this review provides several recommendations for addressing particle size in sediment source fingerprinting research.

**Table 3:** Examples of sediment tracing research using only elemental geochemistry (e.g. major, trace and rare earth elements).


Reference	Country	Area (km <sup>2</sup> )	Elements Analyzed (n)	Particle Size (µm)	Corrections or other approaches to address particle size dynamics	Pre-treatment before analysis	Raw Data Available
Cooper et al., 2015	United Kingdom	5	11	<63	--	Dispersed <sup>a</sup>	--
D'Haen et al., 2013	Turkey	264	18	<63	--	Dispersed <sup>b</sup>	--
Douglas et al., 2003	Australia	22000	50	<10	--	--	--
Haddadchi et al., 2015	Australia	911	41	<10, 10-63, 63-212	--	--	--
Hardy et al., 2010	Canada	12000	55	63-250	Density separation	--	--
Lacey et al., 2015a	Australia	74	37	<10	--	--	--
Lacey and Olley, 2015	Australia	75, 123, 311	23	<10	--	--	--
Olley and Caitcheon 2000	Australia	650000	10	<10	--	Dispersed <sup>c</sup>	Partial <sup>d</sup>
Vale et al., 2016	New Zealand	5870	44	<63	--	--	--

<sup>a</sup> Source samples sonified before sieving

<sup>b</sup> Samples were boiled in distilled water to disperse soil aggregates

<sup>c</sup> Samples sonified before settling in a water column

<sup>d</sup> Raw data is only available for sediment core samples



**Figure 4:** Changes in mineralogy with particle size in sediment from three locations in the Amazon basin from Gibbs (1967) with the red dashed line added to indicate the impact of fractionating the samples at the  $<10\ \mu\text{m}$  particle size.

## 2.4 Mineral Magnetic Properties

Mineral magnetic properties (e.g. magnetic susceptibility, isothermal remanent magnetisation) have also been widely used to investigate sediment provenance (Blake et al., 2004; Palazón et al., 2015a; Pulley et al., 2015b; Walling et al., 1979) (Table 4). The signatures derived from magnetic minerals may be classified as primary (i.e. from parent material prior to weathering) or secondary (i.e. from chemical processes and other processes and effects) (Hatfield, 2014). As the signature is often derived from parent material, mineral magnetic properties can trace sediment derived from different spatial sources (Caitcheon, 1993). Owing to the potential impact of secondary processes, such as anthropogenic inputs and diagenetic processes, mineral magnetic properties may also provide further source discrimination between different erosion processes (Foster et al., 1998; Pulley et al., 2015b).

Magnetic minerals occur in soil and sediment as aggregated concretions, discrete fine grains and particle coatings on very fine grains (Oldfield, 1991; Smith, 1999). Accordingly, mineral magnetic properties may be highly dependent on particle size (Foster et al., 1998; Oldfield et al., 1985). In particular, Hatfield and Maher (2008) demonstrated the importance of characterizing mineral magnetic properties with a particle-size specific approach as different magnetic properties were preferentially associated with different particle size fractions. These authors demonstrated that bacterial magnetosomes formed in lake sediment in the <2  $\mu\text{m}$  and the 2-8  $\mu\text{m}$  fractions. Accordingly, they only quantified source contributions to the 8-31  $\mu\text{m}$  and 31-63  $\mu\text{m}$  fractions. In contrast, Pulley et al. (2015b) reported significantly different magnetic properties in the <32  $\mu\text{m}$  and the >32  $\mu\text{m}$  fractions and, for Caitcheon (1998), the 63-125  $\mu\text{m}$  fraction was appropriate for tracing tributary source contributions.

The challenge is that the particle size fraction driving the mineral magnetic signature may vary for each catchment. For example, the sand and silt fractions may be the most appropriate particle size in catchments in England to quantify sediment sources with magnetic fingerprinting techniques as these fractions have been found to contribute a significant proportion of the mineral magnetic signature (Hatfield and Maher, 2009). Conversely, high magnetic parameters were found in the clay-size particles in the eastern United States (Oldfield et al., 1985). In South Africa, the impact of particle size on mineral magnetic properties was found to be limited in the >32  $\mu\text{m}$  fraction (Pulley et al., 2015b).



**Table 4:** Examples of sediment source fingerprinting research with only mineral magnetic analyses

Reference	Country	Area (km <sup>2</sup> )	Magnetic Parameters (n)	Particle Size (µm)	Corrections or other approaches to address particle size dynamics	Pre-treatment before analysis	Raw Data Available
Blake et al., 2004	Australia	446	9	<10	--	--	Yes
Caitcheon, 1993	Australia	22	2	7 fractions <sup>a</sup>	Density separation	--	--
Foster et al., 1998	United Kingdom	1.5, 12	10	11 fractions <sup>b</sup>	--	Dispersed <sup>c</sup>	--
Hatfield and Mayer, 2009	United Kingdom	240	10	<2, 2-8, 8-31, 31-63, >63	--	Dispersed <sup>d</sup>	--
Oldfield et al., 1985	United States	33	7	10 fractions <sup>e</sup>	--	Dispersed <sup>f</sup>	--
Pulley et al., 2015b	South Africa	148-5751	6	7 fractions <sup>g</sup>	--	Dispersed <sup>h</sup>	--
Slattery et al., 1995	United Kingdom	6	4	<2, 2-16, 16-63	Source correction <sup>i</sup>	Dispersed <sup>j</sup>	--
Walling et al., 1979	United Kingdom	12	5	--	--	--	--

<sup>a</sup> 7 particle size fractions analyzed: <63, 63-125, 125-250, 250-500, 500-1400, 1400-2000 µm

<sup>b</sup> Sediment cores: no fractionation, surface soils: <63 µm, soil cores: <2000 µm, and one bulk sediment samples split into 11 fractions

<sup>c</sup> Calgon before dry-sieving to 4 φ and for <4 φ sodium hexametaphosphate and anhydrous Na<sub>2</sub>CO<sub>3</sub> for the bulk sample split into 11 fractions.

<sup>d</sup> Fractionated samples were dispersed with Calgon prior to sonification

<sup>e</sup> 10 fractions analyzed (1-10 φ)

<sup>f</sup> Dispersed with Calgon

<sup>g</sup> <32, 32-63, 63-125, 125-250, 250-500, 500-1000, 1000-2000µm

<sup>h</sup> Dispersed with sonification

<sup>i</sup> Corrected to a standard particle size distribution consisting of coarse silt (50%), fine silt (40%) and clay (10%)

<sup>j</sup> Dispersal method not provided

The challenge for mineral magnetic properties, and other sediment fingerprints, is that these properties are related to particle size and this relationship varies from catchment to catchment, subcatchment to subcatchment, and even potentially from event to event. Further, these sediment properties are not only sensitive to changes in their sources, they are also sensitive to changes in sediment transport processes that may impact the potential abundance and availability of the different particle size fractions over a range of spatial temporal scales (Hatfield, 2014).

## **2.5 Composite Fingerprinting Approach**

Peart and Walling (1986) advocated for multiple parameters to be used when quantifying sediment source dynamics in order to improve the overall consistency and reliability of source ascription results. The combination of multiple parameters creates a composite fingerprint (Walling et al., 1993) that allows for an increased number of sources to be modelled and is theorized to be more representative of the linkages between sediments and their sources, potentially reducing false matches which were hypothesized to potentially occur with individual tracer properties (Collins et al., 1996). Accordingly, a composite fingerprinting approach has been broadly applied in sediment source fingerprinting research combining several or all of the following: fallout radionuclides, carbon and nitrogen parameters, element geochemistry, mineral magnetism and other parameters, thereby providing significant source discrimination (Collins et al., 1996; Evrard et al., 2013; Navratil et al., 2012; Owens et al., 2000) (Table 5). One challenge with the composite fingerprinting approach is that each of the potential complexities within the particle size – tracer parameter relationship described above is integrated into the expanded composite fingerprint. A second challenge is that it is difficult, if not impossible, to link outputs of statistical-based approaches (e.g. composite fingerprinting) back to a process-based understanding of sediment dynamics.

**Table 5:** Examples of sediment source fingerprinting research using a composite fingerprinting approach incorporating two or more of types of biogeochemical parameters with geochemistry (Geochem), mineral magnetics (Mags), radionuclides (RN), clay mineralogy (Clay min.), diffuse reflectance infrared Fourier transform spectrometry (DRIFTS), X-ray Diffraction (XRD) and other tracers as listed.

Reference	Country	Area (km <sup>2</sup> )	Parameters	Particle Size (µm)	Corrections or other approaches to address particle size dynamics	Pre-treatment before analysis	Raw Data Available
Ben Slimane et al., 2013	Tunisia	2.6	RN, C, N	<2000	SSA	--	--
Ben Slimane et al., 2016	Tunisia	0.6-4	RN, C	<2000	SSA	--	--
Blake et al., 2006	Borneo	<2	Geochem, N	<125	--	--	--
Blake et al., 2012	United Kingdom	1.5	CSSI, Geochem, Mags	<63	SSA <sup>a</sup>	--	--
Caitcheon et al., 2006	Australia	9051	Geochem, RN	<10	--	--	Partial <sup>b</sup>
Collins et al., 1996	United Kingdom	601, 4325	Geochem, RN, C,N	--	SSA	--	--
Devereux et al., 2010	United States	188	Geochem, RN, C	<63	--	--	--
Douglas et al., 2009	Australia	638	Geochem, RN	<10	--	--	--
Douglas et al., 2006a	Australia	144000	Geochem, Clay Min.	<10	--	--	--
Douglas et al., 2006b	Australia	144000	Geochem, Clay Min.	<10	--	--	--
Evrard et al., 2011	France	907	Geochem, RN	<2000	Sc comparison	--	Partial <sup>c</sup>
Evrard et al., 2013	Mexico	3, 9, 12, 630	Geochem, RN, DRIFTS, C, N, $\delta^{13}\text{C}$	<250	d <sub>50</sub> comparison	--	Partial <sup>c</sup>
Hatfield and Mayer, 2008	United Kingdom	240	Geochem, Mags	<2, 2-8, 8-31, 31-63	--	Dispersed <sup>d</sup>	--
Hughes et al., 2009	Australia	6000	Geochem, RN	<10	--	--	--
Koiter et al., 2013b	Canada	74	Geochem, RN	<2000	--	--	--
Le Gall et al., 2016	France	24	Geochem, RN, Sr Isotopes	<63, <2000	Th-correction	--	Yes
Martínez-Carreras et al., 2010	Luxembourg	0.7, 3, 4	Geochem, RN, Colour, C, N, P	<63	--	--	--
Minella et al., 2008	Brazil	1.2	Geochem, C	<150	--	--	--
Motha et al., 2002	Australia	110	Geochem, RN, Mags	<2, 2-20, 20-40, 40-63	Source correction <sup>e</sup>	Dispersed <sup>f</sup>	--
Motha et al., 2003	Australia	65	Geochem, RN	<2, 2-20, 20-40, 40-63	Source correction <sup>e</sup>	Dispersed <sup>f</sup>	--
Mukandan et al., 2010	United States	182	Geochem, RN, $\delta^{15}\text{N}$ , C, N	<2000	Texture comparison	--	--

Navratil et al., 2012	France	905	Geochem, RN	<63	--	--	Partial <sup>c</sup>
Owens et al., 2000	United Kingdom	4390	Geochem, RN, Mags, C, N	<63	SSA	Dispersed <sup>d</sup>	--
Owens et al., 2006	Canada	135, 215	Geochem, Mags, C, N	<500	--	--	--
Palazon et al., 2015a	Spain	1509	Geochem, Mags, RN, C	<63	--	--	Partial <sup>c</sup>
Poleto et al., 2009	Brazil	0.8	Geochem, C	<63	SSA	--	--
Pulley et al., 2015a	United Kingdom	1634	Geochem, Mags, RN	<63	SSA / None	--	--
Russell et al., 2001	United Kingdom	1.5, 4	Geochem, Mags, RN, C, N	<2, 10, 38, 63	Tracer specific particle size correction factor	--	--
Tiecher et al., 2016	Brazil	1.2	Geochem, DRIFTS, XRD	<63	--	--	--
Sherriff et al., 2015	United Kingdom	11	Geochem, Mags	<125	--	--	--
Smith and Blake 2014	United Kingdom	920	Geochem, RN, C	<63	SSA / None	--	--
Stone et al., 2014	Canada	751	Geochem, C	<63	SSA	--	--
Walling et al., 1993	United Kingdom	12, 46	RN, Mags, C,N	<63	None (Mags) and source correction (FRN)	--	--
Walling et al., 1999	United Kingdom	818, 3315	Geochem, Mags, RN,C,N	<63	SSA	--	--
Zhang and Liu, 2016	United States	15.6	Geochem, C,N	<53	--	--	--

<sup>a</sup> SSA correction was only used for the composite fingerprinting approach, not the CSSI tracers

<sup>b</sup> Only geochemistry data is available

<sup>c</sup> Sediments not sources

<sup>d</sup> Dispersed with Calgon and then sonified

<sup>e</sup> Fractional mass of each sediment size fraction was multiplied by their corresponding source tracer property and summed

<sup>f</sup> Dispersed with sonification

### 3. Main Approaches to Address Particle Size

Owing to the potential of particle size to affect the values of tracer properties, researchers have adopted two main approaches to predict or mitigate particle size effects on sediment source fingerprints: fractionation and particle size corrections. Tables 1 to 5 summarize the different approaches used by researchers to address particle size for fallout radionuclides (Table 1), carbon and nitrogen parameters (Table 2), elemental geochemistry (Table 3), mineral magnetic properties (Table 4) and combinations of multiple parameters in a composite fingerprinting approach (Table 5).

#### 3.1 Fractionation

To address particle size impacts on sediment fingerprint properties, researchers often fractionate both their sediment and source samples to a specific and comparable particle size fraction using settling based on Stokes' Law or sieving. The objective is to minimize potential sorting-induced differences between source and sediment properties. Conceptually, any particle size fraction can be isolated to attempt to achieve this objective, though it is mainly the <10  $\mu\text{m}$  and <63  $\mu\text{m}$  fractions that have been isolated in sediment source fingerprinting research.

The <10  $\mu\text{m}$  fraction is predominantly used in Australia to research the source of very fine silt and clay material. The logic supporting the use of the <10  $\mu\text{m}$  fraction is that it is the dominant size fraction being transported in these river systems and this fraction has the greatest ecological and water quality impact (Douglas et al., 2003; Olley and Caitcheon, 2000). The <10  $\mu\text{m}$  fraction is isolated in settling columns based on Stokes' Law with assumptions of constant temperature, roundness and density of the particles (Fontaine et al., 2000; Walden and Slattery, 1993).

The <63  $\mu\text{m}$  fraction is arguably the most adopted sediment tracing particle size fraction (Devereux et al., 2010; Pulley et al., 2015a; Walling et al., 1993). This fraction represents the silt and clay material that is transported preferentially as suspended sediment in riverine, lacustrine, estuarine and coastal systems. The <63  $\mu\text{m}$  fraction is also one of the smallest dry/wet sieve sizes at the fine sand to silt size boundary (i.e. an operationally defined fraction/separation). Research examining the different impacts of wet sieving and dry sieving on biogeochemical properties may be warranted in the sediment source fingerprinting context, along with research on the impact of different pre-treatments to disperse aggregates prior to analyses (Tables 1-5).

One often overlooked component of addressing particle size is that the fraction isolated should include the range of fingerprint property values in the potential sources. This is particularly important for elemental geochemistry. For example in Figure 4, it is apparent that isolating the <2 µm, <10 µm and <63 µm fractions will likely result in different elemental compositions for the three sources based on the particle size fraction selected. This may be particularly important with mineral magnetic tracer properties where different signatures exist in the <32 and the 32-63 µm soil and sediment fractions, indicating that sieving to <63 µm may be inappropriate in some regions (Pulley et al., 2015b).

A second overlooked component is that the particle size fraction isolated for sediment property analyses should directly relate to the research objective. For example, researchers have shown that if the objective is to examine the source of material degrading the Great Barrier Reef near Australia, the <16 µm particle size fraction should likely be targeted (Bartley et al., 2014). Researchers should support their choice of particle size fraction by relating it to the particle size being transported in the stream system (Wallbrink et al., 1999), or the particle size of the sediment-associated contaminants of interest (Olley and Caitcheon, 2000). For both approaches, there is a trade-off between fractionating down to the finest particle size (e.g. <2 µm) versus using a broader particle size fraction (e.g. <63 µm) that may require more steps to address discrepancies between source and sediment particle size distributions. In some regions, the abundance of material present in the sediment and sources may even control the particle size selected as there may not be sufficient <10 µm or even <63 µm material available for analysis. Indeed, the results from sediment source fingerprinting research will only relate to the particle size fraction examined. If narrow particle size ranges are fractionated (e.g. <2 µm or <10 µm), the results from the source apportionment modelling will only apply to that fraction and not the entire suspended sediment load more generally (Mukundan et al., 2012).

Ultimately, the key to using fractionation to address particle size differences is to ensure that the grain size distribution of the source material is similar to that of the sediment sampled (i.e. Poulenard et al., 2009). For example, Sherriff et al. (2015) found that the 90<sup>th</sup> percentile of the suspended sediment distribution for select samples was frequently >63 µm and thus these authors sieved samples to <125 µm. Furthermore, it is important to understand the relationship between particles size distribution and event magnitude, and even how this relationship is impacted by seasonality (Bogen, 1992; Lewis, 1996). In this regard, it would be beneficial for fingerprinting studies to report summary statistics for particle size data of both source soils and sediments to present this important comparison. The challenge is that particle size fractionation of samples alone will not necessarily reduce the discrepancy between the fractionated

samples (Cooper et al., 2015a; Kersten and Smedes, 2002). If there are significant differences between source and sediment particle size distributions remaining after fractionation, particle size corrections are potentially required.


### **3.2 Particle Size Corrections**

To mitigate differences in the particle size distributions of source soil and sediment, corrections have been applied based on particle size characteristics of source and sediment material (Collins et al., 1996; Slattery et al., 1995; Walling and Woodward, 1992; Walling et al., 1993) (Tables 1, 4, 5). The initial corrections reconstructed particle size distribution and tracer parameters of the source materials to allow for the direct comparison with the sampled sediment (Slattery et al., 1995; Walling and Woodward, 1992; Walling et al., 1993). The objective of these corrections was to reduce the impact of source and sediment particle size distributions on tracer parameters prior to quantifying source contributions.

A second approach to particle size corrections is the incorporation of a within-model weighting (Collins et al., 1996) which has been broadly applied (Collins et al., 2012; Poletto et al., 2009; Stone et al., 2014; Walling et al., 1999). The within-model correction incorporates some variant of a particle size weighting, such as the ratio of the mean SSA in sediment to the mean SSA in each source (Collins et al., 1996; Collins et al., 2010). The logic supporting the use of a SSA derived correction is that particle size and specific surface area are closely related (Horowitz, 1991). Surface area is one of the most important controls on sediment trace element concentrations as the majority of these interactions are postulated to be related to surface area chemistry or surface area reactions (Horowitz and Elrick, 1987).

Although widely applied, there are acknowledged limitations and challenges with particle size corrections. Russell et al. (2001) reported a large range in SSA between different sources and sediment for a catchment in the UK and found that a linear-based particle size correction may be inappropriate in some cases. Smith and Blake (2014) further demonstrated that the fundamental assumption of some particle size corrections (i.e. positive linearity between particle size and tracer concentration) does not apply to all tracer properties or equally to properties from different sources (Figure 5). These authors reported that this basic assumption of linearity needs to be constantly examined and its dependence on analytical methods (e.g. acid leached versus total digestion for ICP-MS analyses) should also be considered. Particle size corrections, if adopted, probably should be more rigorous than simple SSA ratio model weightings (e.g. Motha et al., 2002; Motha et al., 2003; Russell et al., 2001; Slattery et al., 1995; Walling and Woodward,

1992). There may even be potential to incorporate particle size properties and organic matter content effects simultaneously with a stepwise multiple regression analysis model (Kraushaar et al., 2015).



**Figure 5:** Pearson correlation coefficients (r) between Specific Surface Area (SSA) and tracer property for three sources from Smith and Blake (2014) (data from Table 1) with the \* indicating fallout radionuclides ( $^{137}\text{Cs}$  and  $^{210}\text{Pb}_{\text{ex}}$ ) on the x axis and the remainder being elements.

The challenge is that there are a variety of non-linear responses between sediment biogeochemistry and particle size (Motha et al., 2002; Russell et al., 2001; Smith and Blake, 2014). There are assumptions with particle size measurements that are often not acknowledged (e.g. all particles are spherical and transported as discrete particles) and there are uncertainties and errors with laser particle size measurements that are not often reported nor propagated into total modelling uncertainty. The impact of these errors varies for different particle size distributions, with a notable increase in instrumentation error with decreasing particle size (Merkus, 2009) (Figure 6). One question for future research is whether or not potential errors on the SSA analysis could result in a substantially different interpretation of the results. Further, stable soil aggregates and agglomerated composite particles that are formed by particle to particle interaction after mobilization (Droppo et al., 2005) may also impact relationships between source and sediment fingerprints. Given the potential uncertainty of accurate SSA ratio-based corrections,



researchers often do not apply particle size corrections and instead rely on physical reasoning of sediment transport processes and/or sample fractionation (e.g. Koiter et al., 2013a; Martinez-Carreras et al., 2010).




Figure 6: Relationship between potential sources of error and particle size distribution, adapted from Merkus (2009).

### 3.3 Combining Size Fractionation and Corrections Procedures

Moving forward, the first step towards addressing particle size should be to fractionate the source and sediment material according to the research question (Table 6), which in turn requires information on the particle size composition of the sediment and/or contaminants in question. Second, the results of this fractionation should be assessed to determine whether the particle size distributions of the source and sediment material are not significantly different. If they are significantly different, third, the application of corrections to tracer properties is probably required for each source with enough samples to ensure source representativeness. Particle size corrections are likely to be more effective if researchers are working with a wide target fraction that is susceptible to sorting effects. The actual impact of the corrections should also be assessed and reported upon, particularly the impact of the corrections on the original basis for source discrimination. The optimal approach for understanding the impact of particle size corrections involves the comparison of biogeochemical properties from samples fractionated across a variety of particle size fractions (e.g. He and Walling, 1996; Laceby et al., 2015b; Russell et al., 2001). Understanding the impacts of particle size on tracer properties should be one of the fundamental first steps when trialling new tracer properties in sediment source fingerprinting research.

**4. Alternative Techniques to Address Particle Size**

Although particle size fractionation and particle size corrections are the two main approaches used to account for any predictable changes in biogeochemical properties during sediment generation, transportation and deposition processes, the challenges with particle size selectivity have opened up new avenues to explore alternative approaches and research directions.

**4.1 Tributary (or Confluence) Tracing**


One approach to mitigating potential particle size impacts on sediment source fingerprinting is to incorporate a tributary tracing or confluence tracing research design (Caitcheon, 1993; Hatfield and Maher, 2008; Laceby et al., 2015a; Olley and Caitcheon, 2000; Vale et al., 2016; Walling et al., 1999). The concept of a tributary tracing approach is that researchers sample sediment in the different upstream tributaries and use these samples as a potential source of sediment sampled further downstream (Figure 7). The tributary sampling approach models sediment as a source and a sink, thus removing a significant proportion of the impact of potential particle size enrichment on fingerprint properties. This approach has also been recently applied to a lacustrine environment by Le Gall et al. (2016) who modelled the source of material sampled in the downstream section of a pond in France based on the geochemical properties of deposited sediment in the inlets of the two main tributaries.

There may be potential particle size enrichment or depletion impacts on fingerprint properties that may occur during transportation and settling processes, although the most significant particle size enrichment typically occurs during the initial stages of mobilization and transportation that often occurs on hillslopes and in ephemeral systems (Stone and Walling, 1997). As material moves into the riverine system, sediment particle size often becomes increasingly uniform and thus fewer differences may be anticipated. Of course, there are always exceptions. For example, Koiter et al. (2015) used a recirculating flume to demonstrate that it was not only distance travelled that impacted particle size, but also channel bed characteristics such as roughness, porosity and inter-gravel flow. Therefore, the tributary tracing technique may have a limited ability to mitigate particle size impacts where tributaries have distinctly different channel bed characteristics and/or where there is a considerable distance between tributary and downstream sampling sites.

**Table 6:** Overview of particle size implications at different steps in the sediment source fingerprinting research process

Step	Particle Size Implications
------	----------------------------

1. Research Objective	<ul style="list-style-type: none"> <li>• Fractionate source and sediment material according to the research question</li> <li>• The particle size fraction analyzed should relate to the dominant particle size being transported in the system or the dominant particle size of the sediment-associated contaminant of interest</li> </ul>
2. Research Design	<ul style="list-style-type: none"> <li>• Consider the potential impacts of particle size on the research design and the different opportunities available to address and investigate particle size impacts on sediment source fingerprinting results</li> <li>• Consider using a tributary tracing sampling design or edge-of-field samplers to mitigate particle size impacts on tracing parameters</li> <li>• Plan and budget particle size analyses for sediment and source materials to understand whether or not there are significant impacts of particle size selectivity on the tracer parameters in the study region</li> </ul>
3. Sample Processing, Analysis, Modelling	<p>A) Fractionation</p> <ul style="list-style-type: none"> <li>• Ensure the particle size distribution of the source material is not significantly different than that of the sediment</li> <li>• Consider the potential impact of fractionation (e.g. sieving/settling) on biogeochemical properties and potential challenges with stable soil aggregates and sediment flocculants</li> </ul> <p>B) Corrections</p> <ul style="list-style-type: none"> <li>• Consider corrections if fractionation does not remove the differences between source and sediment particle size distributions</li> <li>• Always plot and assess the impact of the corrections on tracer parameters and source discrimination</li> </ul> <p>C) Modelling</p> <ul style="list-style-type: none"> <li>• Consider modelling different particle size fractions or comparing results with and without corrections on artificial mixtures to truly understand particle size impacts in the study region</li> </ul>
5. Results	<ul style="list-style-type: none"> <li>• Present the relationship between particle size and the tracer properties of interest</li> <li>• Assess and report on the impact of fractionation and corrections in the results with an emphasis on any potential impacts on the original basis of source discrimination</li> <li>• Present summary statistics for particle size distributions to facilitate comparisons between particle size distributions in source and sediment material</li> </ul>
6. Discussion	<ul style="list-style-type: none"> <li>• Discuss the impact of particle size in relation to other sediment source fingerprinting studies</li> </ul>
7. Supplementary Information	<ul style="list-style-type: none"> <li>• Provide corrected and uncorrected data for parameters used and all particle size data for future use by the research community (including sample coordinates, <math>d_{50}</math>, <math>d_{90}</math>, SSA and pre-treatments used prior to particle size analyses).</li> </ul>



**Figure 7:** An example of a sampling design with the tributary tracing technique where the letters indicate the different tributaries that could be sampled in this theoretical catchment. In this situation, sediment collected at sites c, e and g is compared to sediment collected where it may be able to infer which tributaries (i.e. a, b, d or f) were the main sediment sources.

The tributary tracing technique may also address potential challenges that arise from variations in tracer properties at the sub-catchment or reach scale. For example, it is conceivable that magnetic properties or elemental concentrations may vary significantly at the subcatchment scale. The tributary tracing approach may capitalize on these variations, incorporating them into contrasts between different sources, in this instance, between different tributaries. A question for future research is what should be the benchmark for indicating that the spatial variability in tracer properties has been sufficiently constrained, let alone the heterogeneity in potential fractionation of different particle sizes?

Future research needs to examine the potential of tributary tracing research designs to address particle size enrichment impacts on biogeochemical tracer properties, particularly when comparing sediment sampled instream to material deposited in lakes and reservoirs, and tributaries with different channel bed roughness. Indeed, more research is also required to examine the impact of in-stream sampling approach (e.g. lag deposits, time-integrated samplers, instantaneous samplers, and channel bed-material samplers) on sediment characteristics as different in-stream sampling methods may result in particle size biases that may impact source modelling results. A sampling design with sufficient replication of these different in-stream sampling methods in various catchments around the world may indeed provide further understanding regarding the relationship between particle size and sampling methodology and how the sediment sampling design may impact fingerprint properties and even potentially mixing model results.

#### **4.2 Edge-of-Field Samplers**


An alternative technique that may limit potential particle size effects on biogeochemical tracers during mobilization and initial transportation phases is the installation of edge-of-field samplers (Panuska et al., 2008; Wallbrink and Murray, 1993). Edge-of-field sampling approaches capture sediment after the initial mobilization and transport processes and are an alternative technique for examining whether particle size corrections are required (Evrard et al., 2016), or if they have been applied correctly (Foucher et al., 2015). Conceptually, samplers installed on USLE plots (Brooks et al., 2014; Wischmeier and Smith, 1978) could opportunistically sample material mobilized from hillslopes. Gerlach troughs (Gerlach, 1967) could sample suspended sediment on hillslopes for further analyses after it has been mobilized and transported downslope. Similarly, V-notch weirs combined with automated sediment samplers may also provide an effective approach for sampling hillslope sediments in ephemeral gullies and other ditch type landscape features (Freebairn and Wockner, 1986).

The key is to understand the impact of the initial mobilization and transport processes on both particle size and the fingerprinting parameters of interest. For example, Evrard et al. (2016) demonstrated that there was no significant difference between the fallout radionuclide activity concentrations in surface soils, riverine sediments and sediments sampled with edge-of-field samplers in Laos. Conversely, in a lowland and well-drained agricultural catchment in France, Foucher et al. (2015) sampled material in an ephemeral rill during a rainfall event to confirm both the hyper enrichment of fine particles in overland flow ( $<2\ \mu\text{m}$ ) and also to provide confidence that both the SSA- and Th-based corrections were applied effectively. Further research is required to examine the potential of edge-of-field samplers to improve our understanding of particle size impacts on biogeochemical tracer properties and understand how to apply these samplers for a range of complex source types (e.g. unpaved roads and farm tracks). For example, it would be beneficial to compare the impact of tributary sampling technique, an edge-of-field sampling approach and top soil grab sampling on particle size distributions and to investigate whether or not these sampling approaches mitigate particle size effects on biogeochemical tracers.

#### **4.3 Multiple Fraction Tracing**

Another technique to examine differences and potential particle size effects on conservative behavior is tracing different particle size fractions (Caitcheon, 1998; Haddadchi et al., 2015; Laceby et al., 2016). Although there has been a significant amount of research invested into developing particle size corrections (Collins et al., 1996; Russell et al., 2001; Walling et al., 1993), less frequently have the different fractions been traced and the results from tracing different size fractions been examined.

Caitcheon (1998) reported that source contributions from a sedimentary rock tributary basin varied only by 13% (i.e. from 63% to 76%) when fingerprinting with mineral magnetic properties from different particle size fractions ( $<63\ \mu\text{m}$ ,  $63\text{-}125\ \mu\text{m}$ ,  $125\text{-}250\ \mu\text{m}$  and  $250\text{-}500\ \mu\text{m}$ ). This author reported that the  $63\text{-}125\ \mu\text{m}$  fraction was most likely representative of the bulk material being transported in this particular catchment. Using elemental geochemistry, Haddadchi et al. (2015) traced the fractionated  $<10\ \mu\text{m}$ ,  $10\text{-}63\ \mu\text{m}$  and  $63\text{-}212\ \mu\text{m}$  fractions. These authors found that although the maximum difference was high (33%), there was  $<15\%$  mean absolute difference between modelled source contributions with these fractions (Figure 8). In summary, Haddadchi et al. (2015) indicated that their results highlight the importance of fractionating the particle size most relevant to the management objective of the research.



**Figure 8:** Mean absolute differences (dashed lines) in the source contributions between the <10 μm and the 10 - 63 μm (blue) and between the <10 μm and the 63 - 212 μm fractions (red) for eight sites in Australia plotted with data from Figure 9 in Haddadchi et al. (2015).

Lacey et al. (2016) also compared the impact of tracing different particle size fractions for three catchments in the Fukushima region in Japan. These authors compared the relative model difference from tracing the <63 μm and the <2 mm fractions with TOC, TN,  $\delta^{13}\text{C}$ , and  $\delta^{15}\text{N}$  for 76 sediment samples taken in three different catchments (Figure 3). The authors reported a maximum modelled average difference between the <2 mm and <63 μm fraction of 14%, 11% and 7% for the three catchments, and a mean relative difference of only 6% (SD 3%) for all possible modelled source contributions. The limited differences between these models was likely driven by the fact that only 19% (SD 13%) of the material in these sediment samples was <63 μm compared to 23% (SD 12%) of the material for the source samples, indicating that ~80% of material was >63 μm for both source and sediment samples. A similar approach should be applied in catchments with finer sediment transiting the system to characterize the potential impact of fine sediment material on modelling results when tracing different fractions.

These studies raise interesting questions regarding what are the actual differences in mixing model results arising from the impacts of particle size on tracer properties. Are these impacts greater than analytical and model uncertainty? Is this particle size impact significant for management objectives? These fundamental

research questions present an opportunity for sediment source fingerprinting research with a focus on methodological sensitivity to particle size effects in different systems. The challenge moving forward will be for researchers to quantify the solid discharge in each particle size fraction in order to compare and model both the relative source contributions (i.e. unweighted) and the absolute source contributions (i.e. weighted by the abundance in each fraction), which could be significantly different.

## **5. Particle Size Opportunities**

After material is mobilized from sources, sediment fingerprinting researchers often conceptualize riverine systems as a black box, where the processes that occur between source and sediment sampling are not well understood (i.e. Figure 3 in Koiter et al., 2013b). If researchers focus on furthering our understanding of the relationship between particle size and biogeochemical properties, they may simultaneously start to examine processes occurring within this black box. De-convoluting the complex relationship between particle size and biogeochemical properties may provide significant insight into the processes of sediment mobilization, transport and deposition within this black box, including the potential unique behaviour of different sediment source parameters. Accordingly, there are multiple opportunities to advance sediment source fingerprinting research and further our understanding of the complex relationship between particle size and tracer parameter predictability.

### **5.1 Mineralogy, Particle Size and Elemental Geochemistry**

The most important research objective and opportunity regarding particle size is to increase our understanding of sediment tracer predictability. One approach to improving this predictability is connecting particle size, mineralogy and elemental geochemistry. For example, in Figure 4, there is a clear connection between sediment mineralogy and particle size, which will have direct ramifications for the elemental compositions of sediment sources. Significant advances in sediment source fingerprinting research are likely to be made through connecting these three components. More research into sediment mineralogy may result in novel tracing techniques that are grounded in a logical basis for source discrimination (e.g. Afshar et al., 2016; Bainbridge et al., 2016; Gingele and De Deckker, 2004; Tiecher et al., 2016), that also provide significant information on the complex relationship between particle size and elemental concentrations.

Mineralogy was one of the first sediment fingerprinting properties (Klages and Hsieh, 1975; Wall and Wilding, 1975) and returning to incorporate mineralogy more directly into sediment tracing may provide significant insights into particle size impacts on elemental compositions. This may be achieved, although



indirectly, through using Diffuse Reflectance Infrared Fourier Transform Spectrometry (DRIFTS) to identify mineralogical groups (e.g. calcite, aluminosilicates, quartz) and discriminate between sediment lithological sources (Poulenard et al., 2012). The use of DRIFTS and other spectroscopy approaches to tracing sediment sources has the potential to develop a strong link to mineralogy and potentially estimate other conventional tracer parameters like texture (Balsam and Deaton, 1996; Ortiz et al., 2009). Other approaches such as clay mineral ratios (e.g. illite/illite + expandable clays) have also shown promise to trace sediment provenance (Bainbridge et al., 2016; Douglas et al., 2006a; Douglas et al., 2006b). Connecting clay mineralogy to element geochemistry and particle size will likely provide complementary information relating to each of these fundamental sediment property characteristics.

## **5.2 Elemental Ratios and Elemental Normalisation**

Moving a step beyond incorporating mineralogy more effectively into sediment tracing research is the potential for incorporating approaches from other disciplines to address particle size effects. For example, in the sediment provenance literature, elemental ratios have long been used to investigate changes in particle size. In particular, the Al/Si ratio is used as a proxy for particle size in sediment provenance research (Figure 9) (Bouchez et al., 2011). Other ratios may indeed be effective, such as the Ca/Al ratio (Chen et al., 2014), along with particle size sensitive elements such as Ti (Bábek et al., 2015), or even simply clay content (Szava-Kovats, 2008). For example, studies reconstructing the evolution of metal enrichment ratios in sediment have also normalized elemental concentrations to Sc, Al, Si, Li or Th to minimize particle size impacts when examining contamination trends (Ayrault et al., 2010; Clark et al., 2014; Grosbois et al., 2012; Kersten and Smedes, 2002; Le Cloarec et al., 2011). Indeed, there are multiple approaches from the sediment provenance field (Armstrong-Altrin et al., 2015; Bábek et al., 2015; Owens et al., 2016; Singh et al., 2005) that present significant opportunities for enhancing the sediment source fingerprinting technique (e.g. Vale et al., 2016).


Different elemental and lithogenic radionuclide ratios have been used previously to address particle size and density related enrichment in sediment source fingerprinting research. For example, Olley and Murray (1994) demonstrated that although the concentrations of thorium (Th) isotopes varied with particle size, the  $^{230}\text{Th}/^{232}\text{Th}$  ratio remains constant, with sorting by density or particle size producing the same ratio as the bulk soil. Caitcheon (1998) and Murray et al. (1993a) incorporated ratios directly into the mixing model process for mineral magnetism and radionuclides, respectively. Although these modelling approaches used the actual ratio lines to quantify source contributions, they demonstrate the utility of incorporating ratios directly into the mixing model process to address particle size enrichment. More research is required to

examine whether similar approaches may also provide a technique for normalizing particle size effects with frequentist or Bayesian mixing models, or even more insight into the foundation for elemental discrimination between sediment sources. For example, the interrelationships between particle size and biogeochemical tracer properties indicate that particle size may directly contribute to the formation of some contrasts in sediment fingerprints between sources. Accordingly, further advances in understanding these relationships, with experimental designs incorporating different particle size fractions, may provide additional understanding for the basis of discrimination between sediment sources.

### **5.3 Particle Dynamics**

The nature of cohesive sediment transport, and particularly fine sediment transport, is fundamentally complex where the majority of particles are transported as flocs, or composite particles, that are comprised of organic matter, smaller particles and mineral components with a complex structure (Droppo et al., 2005; Walling and Collins, 2016). The presence of these soil aggregates and sediment flocs will have a significant impact on particle size selectivity during sediment mobilization, transportation and deposition processes (Beuselinck et al., 2000). The behaviour of material during these processes is a function of density, aggregate stability, grain shape and grain size (D'Haen et al., 2012).

The impact of grain abrasion and disaggregation on  $^{137}\text{Cs}$  concentrations was examined by Dyer and Olley (1999) who reported that the fraction produced by abrasion had essentially similar  $^{137}\text{Cs}$  concentrations as the corresponding fraction ( $<40\text{ }\mu\text{m}$ ) in non-disturbed soil. Conversely, Crockford and Olley (1998) found that breakage and abrasion processes had a substantial impact on mineral magnetic properties, reducing concentrations in a granitic soil and increasing concentrations in a sedimentary soil. These authors concluded that a tributary or confluence tracing approach should mitigate the impact of these processes on mineral magnetic properties. Although these studies examined the disaggregation of sediments, Droppo et al. (2005) suggested that the actual nature of aggregates and flocs is rarely considered in studies of sediment transport and sources.



**Figure 9:** Scatter plots of the relationship between the Al/Si ratio and  $D_{90}$ , K, and Cs (ppm) from samples from different tributaries of the Amazon catchment with data from Tables 1 and 2 from Bouchez et al., (2011).

Accordingly, there is an opportunity to improve our understanding of sediment transport processes by further incorporating the nature of the material being transported more directly into the methodology of sediment fingerprinting projects. This is important as the sediment fractions isolated by density may be

different than those identified by particle size analysis (Gregorich et al., 2006). Different fractionation methods (e.g. wet sieving, dry sieving, grinding) may fragment aggregates which could result in different biogeochemical properties in analyzed aliquots. This may be particularly true with respect to different techniques and conditions for wet and dry sieving, and the use of Stokes' law and settling columns to isolate different particle size fractions. Different source and sediment material (e.g. platy, spherical, and rod-shaped grains) may respond differently to fractionation methods (Droppo et al., 2005; Hatfield, 2014). Different pre-analysis treatments (e.g. sonification, chemical dispersion, physical dispersion) may also impact the biogeochemical fingerprints of interest. Laboratory determined grain size distributions will most likely differ significantly from those in the field owing to the physical and often chemical disaggregation of flocs and aggregates during the laboratory fractionation process (Phillips and Walling, 1995; Walling and Collins, 2016).

One major research opportunity is whether these alterations between laboratory and field, and the impact of aggregates, are significant to sediment fingerprinting modelling results, or fall within the range of analytical and modelling uncertainty. For example, if soil aggregates and sediment flocs form and/or evolve during sediment mobilization and transportation processes, this may result in a form of non-conservative behaviour between source and downstream sink, adding uncertainty in the form of an 'aggregate size' effect on tracer properties. Furthermore, the impact of aggregates directly relates to the approach to sample processing, particularly whether aggregates should be chemically dispersed prior to particle size fractionation or whether samples should not be disaggregated (Koiter et al., 2017). More research is required to characterize the impact of aggregates, and their processing, on sediment source fingerprinting research.

Density separation is an intriguing fractionation approach that has not received much attention in the sediment source fingerprinting literature (Hardy et al., 2010). Different clay minerals have different densities that may impact their separation with settling columns in the finer fractions. Furthermore, iron oxides and mineral magnetic properties have been strongly related to density (Hatfield, 2014) and there may be micro aggregates or sediment flocs transported in suspension with the silt or very fine sand fraction because they are lighter. Importantly, these micro-aggregates and sediment flocs will have a high capacity to transport contaminants; however they will have a different resistance (i.e. a lower density) relative to absolute particles of a similar size, predominantly the silt fraction (Droppo et al., 1998).

Although density separation is time consuming and expensive, it may be more relevant to fluvial processes compared to sieving the <63  $\mu\text{m}$  fraction or settling the <10  $\mu\text{m}$  fraction. Accordingly, it would be

opportunistic to compare density and particle size separation techniques to understand their impacts on sediment fingerprinting modelling results. As researchers apply tracing techniques directly to quantify the relative source contributions of carbon and nitrogen (Cooper et al., 2015b; Garzon-Garcia et al., 2017), and are using different carbon components to model sediment sources (Blake et al., 2012; Gibbs, 2008; Hancock and Revill, 2013; Reiffarth et al., 2016) the utility of density separation may become of increasing importance for tracing different carbon sources in riverine systems.

#### **5.4 Particle Size as a Tracer Property**

Situations may arise where particle size may be used as a tracer property in and of itself. The sediment source fingerprinting approach is founded upon the assumptions of conservative behavior and source discrimination. As noted in the introduction and throughout this review, if the sediment source properties, including particle size, are predictable, they could possibly be used in sediment source fingerprinting research.

The particle size distribution of source material is dependent on the parent material, weathering and erosion processes. Accordingly, there may be situations where particle size itself may be a useful tracer property, particularly where one sediment source is dominated by sandy quartz material and another by fine clay material. Researchers have effectively utilized particle size to discriminate between pre- and post-dam stratigraphy (Batuca and Jordaan Jr, 2000; Morris and Fan, 1998). Further, the particle size of reservoir sediment has been traced back to the parent rock material (Abraham et al., 1999). If researchers were interested in tracing material in the bed load, or material being transported off shore to oceanic environments, then particle size metrics may be effective tracers in these scenarios, if they are indeed predictable. There may even indeed be situations where particle shape may provide an effective discriminator for determining sediment provenance (Ehrlich et al., 1980).

#### **6. Conclusion**

Sediment source fingerprinting researchers have recently tended to avoid in-depth examinations of fundamental topics such as the impact of organic matter on biogeochemical properties, which tracer properties are non-conservative (e.g. soluble, reactive), and what is the impact of particle size on tracer property predictability. One limitation of statistically-oriented approaches to sediment source fingerprinting is that they do not strive to understand the logic of tracer selection, non-conservative tracers, nor particle size impacts. This is particularly limiting, as addressing conservative behaviour is fundamentally more complex than simply acknowledging that sediment samples plot within their source

range. Modelling, and the statistical selection of which tracer properties to model, represent only two stages of the sediment source fingerprinting process; although these two steps, at times, appear to dominate the recent literature. Rarely is particle size the focus of research, rather particle size is often simply an acknowledged limitation that must be addressed somehow. After demonstrating that non-conservative tracer properties affect the accuracy of sediment source fingerprinting modelling results, Sherriff et al. (2015) appropriately emphasized that improved strategies to detect non-conservative tracer properties should be a priority of sediment source fingerprinting research.

Researchers thus need to start investigating some of these fundamental assumptions and complexity at the core of the sediment source fingerprinting approach. Researchers should strive to understand particle size impacts and the logic of tracer selection, as they are likely to be inherently related. For example, the interrelationships between particle size and biogeochemical tracer properties suggest that in many instances particle size may also contribute to the formation of contrasts in sediment fingerprints between sources. Approaches to address particle size will likely vary from region to region, catchment to catchment, and even from event to event. Accordingly, researchers should always publish particle size and biogeochemical property datasets for important regional and future comparisons, including data from different particle size fractions (Kersten and Smedes, 2002). Of note, from the papers reviewed in this study, only approximately 15% published the raw source and sediment data (Tables 1-5).

In summary, particle size dynamics presents both a challenge and an opportunity for sediment source fingerprinting research. The research question, context and objective will determine the approach to addressing particle size and the appropriate fraction for investigation (Table 6). Addressing particle size effects is fundamental to research design, tracer selection, and sampling technique (e.g. edge-of-field samplers). Advances in the understanding of particle size–biogeochemical tracer interactions will likely improve the predictability and therefore accuracy of sediment source fingerprinting.

#### **Acknowledgements**

This research was the result of multiple discussions at European Geosciences Union annual meetings.

698 **References**

- 699 Abraham, J., Allen, P., Dunbar, J. and Dworkin, S., 1999. Sediment type distribution in reservoirs: sediment  
700 source versus morphometry. *Environmental Geology* 38, 101-110.
- 701 Afshar, S., Saghafian, B. and Vosoughifar, H., 2016. Source apportionment of the sediments entering dam  
702 using lithological and mineralogical studies. *Global Journal of Environmental Science and*  
703 *Management* 2, 265-274.
- 704 Amundson, R., Austin, A.T., Schuur, E.A.G., Yoo, K., Matzek, V., Kendall, C., Uebersax, A., Brenner, D. and  
705 Baisden, W.T., 2003. Global patterns of the isotopic composition of soil and plant nitrogen. *Global*  
706 *Biogeochemical Cycles* 17, 1031.
- 707 Armstrong-Altrin, J.S., Machain-Castillo, M.L., Rosales-Hoz, L., Carranza-Edwards, A., Sanchez-Cabeza, J.-A.  
708 and Ruíz-Fernández, A.C., 2015. Provenance and depositional history of continental slope  
709 sediments in the Southwestern Gulf of Mexico unraveled by geochemical analysis. *Continental*  
710 *Shelf Research* 95, 15-26.
- 711 Ayrault, S., Priadi, C.R., Evrard, O., Lefèvre, I. and Bonté, P., 2010. Silver and thallium historical trends in  
712 the Seine River basin. *Journal of Environmental Monitoring* 12, 2177-2185.
- 713 Bábek, O., Grygar, T.M., Faměra, M., Hron, K., Nováková, T. and Sedláček, J., 2015. Geochemical  
714 background in polluted river sediments: How to separate the effects of sediment provenance and  
715 grain size with statistical rigour? *CATENA* 135, 240-253.
- 716 Bainbridge, Z., Lewis, S., Smithers, S., Wilkinson, S., Douglas, G., Hillier, S. and Brodie, J., 2016. Clay mineral  
717 source tracing and characterisation of Burdekin River (NE Australia) and flood plume fine  
718 sediment. *Journal of Soils and Sediments* 16, 687-706.
- 719 Balesdent, J., Mariotti, A. and Guillet, B., 1987. Natural  $^{13}\text{C}$  abundance as a tracer for studies of soil organic  
720 matter dynamics. *Soil Biology and Biochemistry* 19, 25-30.
- 721 Balsam, W.L. and Deaton, B.C., 1996. Determining the composition of late Quaternary marine sediments  
722 from NUV, VIS, and NIR diffuse reflectance spectra. *Marine Geology* 134, 31-55.
- 723 Bartley, R., Bainbridge, Z.T., Lewis, S.E., Kroon, F.J., Wilkinson, S.N., Brodie, J.E. and Silburn, D.M., 2014.  
724 Relating sediment impacts on coral reefs to watershed sources, processes and management: A  
725 review. *Science of The Total Environment* 468–469, 1138-1153.
- 726 Batuca, D.G. and Jordaan Jr, J., 2000. *Silting and Desilting of Reservoirs*. A.A. Balkema Publishers,  
727 Rotterdam, 357 pp.
- 728 Bellanger, B., Huon, S., Velasquez, F., Valles, V., Girardin, C. and Mariotti, A., 2004. Monitoring soil organic  
729 carbon erosion with  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  on experimental field plots in the Venezuelan Andes. *CATENA*  
730 58, 125-150.
- 731 Belmont, P., Gran, K.B., Schottler, S.P., Wilcock, P.R., Day, S.S., Jennings, C., Lauer, J.W., Viparelli, E.,  
732 Willenbring, J.K., Engstrom, D.R. and Parker, G., 2011. Large shift in source of fine sediment in the  
733 Upper Mississippi River. *Environmental Science and Technology* 45, 8804–8810.
- 734 Belmont, P., Willenbring, J., Schottler, S., Marquard, J., Kumarasamy, K. and Hemmis, J., 2014. Toward  
735 generalizable sediment fingerprinting with tracers that are conservative and nonconservative over  
736 sediment routing timescales. *Journal of Soils and Sediments* 14, 1479-1492.
- 737 Ben Slimane, A., Raclot, D., Evrard, O., Sanaa, M., Lefèvre, I., Ahmadi, M., Tounsi, M., Rumpel, C., Mammou,  
738 A.B. and Le Bissonnais, Y., 2013. Fingerprinting sediment sources in the outlet reservoir of a hilly  
739 cultivated catchment in Tunisia. *Journal of Soils and Sediments* 13, 801-815.
- 740 Ben Slimane, A., Raclot, D., Evrard, O., Sanaa, M., Lefevre, I. and Le Bissonnais, Y., 2016. Relative  
741 contribution of rill/interrill and gully/channel erosion to small reservoir siltation in Mediterranean  
742 environments. *Land Degradation and Development* 27, 785-797.
- 743 Beuselinck, L., Govers, G., Steegen, A. and Poesen, J., 2000. Impact of soil aggregates on the size selectivity  
744 of the sediment deposition process. In: M. Stone (Editor), *The Role of Erosion and Sediment*

745 Transport in Nutrient and Contamination Transfer, IAHS Publication No. 263, IAHS Press,  
746 Wallingford, 99-107.

747 Blake, W., Walsh, R., Sayer, A. and Bidin, K., 2006. Quantifying fine-sediment sources in primary and  
748 selectively logged rainforest catchments using geochemical tracers. *Water, Air, & Soil Pollution:*  
749 *Focus* 6, 615-623.

750 Blake, W.H., Ficken, K.J., Taylor, P., Russell, M.A. and Walling, D.E., 2012. Tracing crop-specific sediment  
751 sources in agricultural catchments. *Geomorphology* 139–140, 322-329.

752 Blake, W.H., Wallbrink, P.J., Doerr, S.H., Shakesby, R.A. and Humphreys, G.S., 2004. Sediment  
753 redistribution following wildfire in the Sydney region, Australia: a mineral magnetic tracing  
754 approach. In: V. Golosov, V.R. Belyaev and D.E. Walling (Editors), *Sediment Transfer through the*  
755 *Fluvial System*, IAHS Publication No. 288, IAHS Press, Wallingford, 52-59.

756 Blake, W.H., Wallbrink, P.J., Wilkinson, S.N., Humphreys, G.S., Doerr, S.H., Shakesby, R.A. and Tomkins,  
757 K.M., 2009. Deriving hillslope sediment budgets in wildfire-affected forests using fallout  
758 radionuclide tracers. *Geomorphology* 104, 105-116.

759 Bogen, J., 1992. Monitoring grain size of suspended sediments in rivers. In: J. Bogen, D.E. Walling and T.  
760 Day (Editors), *Erosion and Sediment Transport Monitoring Programmes in River Basins*, IAHS  
761 Publication No. 210, IAHS Press, Wallingford, 183-190.

762 Bonniwell, E.C., Matisoff, G. and Whiting, P.J., 1999. Determining the times and distances of particle transit  
763 in a mountain stream using fallout radionuclides. *Geomorphology* 27, 75-92.

764 Bouchez, J., Gaillardet, J., France-Lanord, C., Maurice, L. and Dutra-Maia, P., 2011. Grain size control of  
765 river suspended sediment geochemistry: Clues from Amazon River depth profiles. *Geochemistry,*  
766 *Geophysics, Geosystems* 12, 1-24.

767 Bradford, J., Truman, C. and Huang, C., 1992. Comparison of three measures of resistance of soil surface  
768 seals to raindrop splash. *Soil Technology* 5, 47-56.

769 Brooks, A., Spencer, J., Borombovits, D., Pietsch, T. and Olley, J., 2014. Measured hillslope erosion rates in  
770 the wet-dry tropics of Cape York, northern Australia: Part 2, RUSLE-based modeling significantly  
771 over-predicts hillslope sediment production. *CATENA* 122, 1-17.

772 Caitcheon, G.G., 1993. Sediment Source Tracing Using Environmental Magnetism - a New Approach with  
773 Examples from Australia. *Hydrological Processes* 7, 349-358.

774 Caitcheon, G.G., 1998. The significance of various sediment magnetic mineral fractions for tracing  
775 sediment sources in Killimicat Creek. *CATENA* 32, 131-142.

776 Caitcheon, G.G., Douglas, G. and Palmer, M., 2006. Sediment Source Tracing in the Lake Burraborang  
777 Catchment. *CSIRO Land and Water*, Canberra, 28.

778 Chen, J.B., Gaillardet, J., Bouchez, J., Louvat, P. and Wang, Y.N., 2014. Anthropophile elements in river  
779 sediments: Overview from the Seine River, France. *Geochemistry, Geophysics, Geosystems* 15,  
780 4526-4546.

781 Clark, D.E., Vogels, M.F., van der Perk, M., Owens, P.N. and Petticrew, E.L., 2014. Effects of a small-scale,  
782 abandoned gold mine on the geochemistry of fine stream-bed and floodplain sediments in the  
783 Horsefly River watershed, British Columbia, Canada. *Mineralogical Magazine* 78, 1491-1504.

784 Clark, E.H., 1985. The off-site costs of soil erosion. *Journal of Soil and Water Conservation* 40, 19-22.

785 Collins, A.L. and Walling, D.E., 2002. Selecting fingerprint properties for discriminating potential suspended  
786 sediment sources in river basins. *Journal of Hydrology* 261, 218-244.

787 Collins, A.L., Walling, D.E. and Leeks, G.J.L., 1996. Composite fingerprinting of the spatial source of fluvial  
788 suspended sediment: a case study of the Exe and Severn river basins, United Kingdom.  
789 *Géomorphologie: Relief, Processus, Environnement* 2, 41-53.

790 Collins, A.L., Walling, D.E., Webb, L. and King, P., 2010. Apportioning catchment scale sediment sources  
791 using a modified composite fingerprinting technique incorporating property weightings and prior  
792 information. *Geoderma* 155, 249-261.



- Collins, A.L., Zhang, Y., McChesney, D., Walling, D.E., Haley, S.M. and Smith, P., 2012. Sediment source tracing in a lowland agricultural catchment in southern England using a modified procedure combining statistical analysis and numerical modelling. *Science of The Total Environment* 414, 301-317.
- Cooper, R.J., Krueger, T., Hiscock, K.M. and Rawlins, B.G., 2014. Sensitivity of fluvial sediment source apportionment to mixing model assumptions: A Bayesian model comparison. *Water Resources Research* 50, 9031-9047.
- Cooper, R.J., Krueger, T., Hiscock, K.M. and Rawlins, B.G., 2015a. High-temporal resolution fluvial sediment source fingerprinting with uncertainty: a Bayesian approach. *Earth Surface Processes and Landforms* 40, 78-92.
- Cooper, R.J., Pedentchouk, N., Hiscock, K.M., Disdler, P., Krueger, T. and Rawlins, B.G., 2015b. Apportioning sources of organic matter in streambed sediments: An integrated molecular and compound-specific stable isotope approach. *Science of The Total Environment* 520, 187-197.
- Crockford, R. and Olley, J., 1998. The effects of particle breakage and abrasion on the magnetic properties of two soils. *Hydrological Processes* 12, 1495-1505.
- D'Haen, K., Verstraeten, G., Duser, B., Degryse, P., Haex, J. and Waelkens, M., 2013. Unravelling changing sediment sources in a Mediterranean mountain catchment: a Bayesian fingerprinting approach. *Hydrological Processes* 27, 896-910.
- D'Haen, K., Verstraeten, G. and Degryse, P., 2012. Fingerprinting historical fluvial sediment fluxes. *Progress in Physical Geography* 36, 154-186.
- Davis, C.M. and Fox, J.F., 2009. Sediment fingerprinting: Review of the method and future improvements for allocating nonpoint source pollution. *Journal of Environmental Engineering* 135, 490-504.
- Devereux, O.H., Prestegard, K.L., Needelman, B.A. and Gellis, A.C., 2010. Suspended-sediment sources in an urban watershed, Northeast Branch Anacostia River, Maryland. *Hydrological Processes* 24, 1391-1403.
- Douglas, G., Caitcheon, G. and Palmer, M., 2009. Sediment source identification and residence times in the Maroochy River estuary, southeast Queensland, Australia. *Environmental Geology* 57, 629-639.
- Douglas, G., Ford, P., Palmer, M., Noble, R. and Packett, R., 2006a. Fitzroy River Basin, Queensland, Australia. I. Identification of sediment sources in impoundments and flood events. *Environmental Chemistry* 3, 364-376.
- Douglas, G., Ford, P., Palmer, M., Noble, R. and Packett, R., 2006b. Fitzroy River, Queensland, Australia. II. Identification of sources of estuary bottom sediments. *Environmental Chemistry* 3, 377-385.
- Douglas, G., Palmer, M. and Caitcheon, G., 2003. The provenance of sediments in Moreton Bay, Australia: a synthesis of major, trace element and Sr-Nd-Pb isotopic geochemistry, modelling and landscape analysis. *Hydrobiologia* 494, 145-152.
- Droppo, I., Nackaerts, K., Walling, D. and Williams, N., 2005. Can flocs and water stable soil aggregates be differentiated within fluvial systems? *CATENA* 60, 1-18.
- Droppo, I., Walling, D. and Ongley, E., 1998. Suspended sediment structure: implications for sediment and contaminant transport modelling. In: W. Summer, E. Klaghofer and W. Zhang (Editors), *Modelling Soil Erosion, Sediment Transport and Closely Related Hydrological Processes*, IAHS Publication No. 249, IAHS Press, Wallingford, 437-444.
- Dyer, F.J. and Olley, J.M., 1999. The effects of grain abrasion and disaggregation on <sup>137</sup>Cs concentrations in different size fractions of soils developed on three different rock types. *CATENA* 36, 143-151.
- Ehrlich, R., Brown, P.J., Yarus, J.M. and Przygocki, R.S., 1980. The origin of shape frequency distributions and the relationship between size and shape. *Journal of Sedimentary Research* 50, 475-483.

839 Evrard, O., Laceby, J.P., Huon, S., Lefèvre, I., Sengtaheuanghoung, O. and Ribolzi, O., 2016. Combining  
840 multiple fallout radionuclides ( $^{137}\text{Cs}$ ,  $^7\text{Be}$ ,  $^{210}\text{Pb}_{\text{xs}}$ ) to investigate temporal sediment source  
841 dynamics in tropical, ephemeral riverine systems. *Journal of Soils and Sediments* 16, 1130-1144.  
842 Evrard, O., Navratil, O., Ayrault, S., Ahmadi, M., Némery, J., Legout, C., Lefèvre, I., Poirel, A., Bonté, P. and  
843 Esteves, M., 2011. Combining suspended sediment monitoring and fingerprinting to determine  
844 the spatial origin of fine sediment in a mountainous river catchment. *Earth Surface Processes and  
845 Landforms* 36, 1072-1089.  
846 Evrard, O., Némery, J., Gratiot, N., Duvert, C., Ayrault, S., Lefèvre, I., Poulenard, J., Prat, C., Bonté, P. and  
847 Esteves, M., 2010. Sediment dynamics during the rainy season in tropical highland catchments of  
848 central Mexico using fallout radionuclides. *Geomorphology* 124, 42-54.  
849 Evrard, O., Poulenard, J., Némery, J., Ayrault, S., Gratiot, N., Duvert, C., Prat, C., Lefèvre, I., Bonté, P. and  
850 Esteves, M., 2013. Tracing sediment sources in a tropical highland catchment of central Mexico by  
851 using conventional and alternative fingerprinting methods. *Hydrological Processes* 27, 911-922.  
852 Fan, Q., Yamaguchi, N., Tanaka, M., Tsukada, H. and Takahashi, Y., 2014. Relationship between the  
853 adsorption species of cesium and radiocesium interception potential in soils and minerals: an  
854 EXAFS study. *Journal of Environmental Radioactivity* 138, 92-100.  
855 Fontaine, T.A., Moore, T.D. and Burgoa, B., 2000. Distributions of contaminant concentration and particle  
856 size in fluvial sediment. *Water Research* 34, 3473-3477.  
857 Foster, I.D.L., Lees, J.A., Owens, P.N. and Walling, D.E., 1998. Mineral magnetic characterization of  
858 sediment sources from an analysis of lake and floodplain sediments in the catchments of the Old  
859 Mill Reservoir and Slapton Ley, South Devon, UK. *Earth Surface Processes and Landforms* 23, 685-  
860 703.  
861 Foucher, A., Laceby, J.P., Salvador-Blanes, S., Evrard, O., Le Gall, M., Lefèvre, I., Cerdan, O., Rajkumar, V.  
862 and Desmet, M., 2015. Quantifying the dominant sources of sediment in a drained lowland  
863 agricultural catchment: The application of a thorium-based particle size correction in sediment  
864 fingerprinting. *Geomorphology* 250, 271-281.  
865 Fox, J.F. and Papanicolaou, A.N., 2007. The use of carbon and nitrogen isotopes to study watershed erosion  
866 processes. *Journal of the American Water Resources Association* 43, 1047-1064.  
867 Freebairn, D. and Wockner, G., 1986. A study of soil erosion on vertisols of the eastern Darling Downs,  
868 Queensland .II. The effect of soil, rainfall, and flow conditions on suspended sediment losses. *Soil  
869 Research* 24, 159-172.  
870 Frings, R.M., 2008. Downstream fining in large sand-bed rivers. *Earth-Science Reviews* 87, 39-60.  
871 Fry, B., 2006. *Stable Isotope Ecology*. Springer. New York., 308 pp.  
872 Garzon-Garcia, A., Laceby, J.P., Olley, J.M. and Bunn, S.E., 2017. Differentiating the sources of fine  
873 sediment, organic matter and nitrogen in a subtropical Australian catchment. *Science of The Total  
874 Environment* 575, 1384–1394.  
875 Gateuille, D., Evrard, O., Lefevre, I., Moreau-Guigon, E., Alliot, F., Chevreuil, M. and Mouchel, J.-M., 2014.  
876 Mass balance and decontamination times of Polycyclic Aromatic Hydrocarbons in rural nested  
877 catchments of an early industrialized region (Seine River basin, France). *Science of The Total  
878 Environment* 470, 608-617.  
879 Gerlach, T., 1967. Hillslope troughs for measuring sediment movement. *Revue de Geomorphologie  
880 Dynamique* 17, 173.  
881 Gibbs, M., 2008. Identifying source soils in contemporary estuarine sediments: A new compound-specific  
882 isotope method. *Estuaries and Coasts* 31, 344-359.  
883 Gibbs, R.J., 1967. Amazon River: Environmental Factors That Control Its Dissolved and Suspended Load.  
884 *Science* 156, 1734-1737.  
885 Gibbs, R.J., Matthews, M.D. and Link, D.A., 1971. The relationship between sphere size and settling  
886 velocity. *Journal of Sedimentary Research* 41, 7-18.

- Gingele, F. and De Deckker, P., 2004. Fingerprinting Australia's rivers with clay minerals and the application for the marine record of climate change. *Australian Journal of Earth Sciences* 51, 339-348.
- Gourdin, E., Evrard, O., Huon, S., Reyss, J.-L., Ribolzi, O., Bariac, T., Sengtaheuanghoung, O., Ayrault, S. and A.L., C., 2014. Spatial and temporal variability of  $^7\text{Be}$  and  $^{210}\text{Pb}$  wet deposition during four successive monsoon storms in a catchment of northern Laos. *Journal of Environmental Radioactivity* 136, 195-205.
- Gregorich, E., Beare, M., McKim, U. and Skjemstad, J., 2006. Chemical and biological characteristics of physically uncomplexed organic matter. *Soil Science Society of America Journal* 70, 975-985.
- Grosbois, C., Meybeck, M., Lestel, L., Lefèvre, I. and Moatar, F., 2012. Severe and contrasted polymetallic contamination patterns (1900–2009) in the Loire River sediments (France). *Science of The Total Environment* 435, 290-305.
- Guzmán, G., Quinton, J., Nearing, M., Mabit, L. and Gómez, J., 2013. Sediment tracers in water erosion studies: current approaches and challenges. *Journal of Soils and Sediments* 13, 816-833.
- Haddadchi, A., Olley, J. and Laceby, J.P., 2014. Accuracy of mixing models in predicting sediment source contributions. *Science of The Total Environment* 497–498, 139-152.
- Haddadchi, A., Olley, J. and Pietsch, T., 2015. Quantifying sources of suspended sediment in three size fractions. *Journal of Soils and Sediments* 15, 2086-2100.
- Haddadchi, A., Ryder, D.S., Evrard, O. and Olley, J., 2013. Sediment fingerprinting in fluvial systems: review of tracers, sediment sources and mixing models. *International Journal of Sediment Research* 28, 560-578.
- Hancock, G.J. and Revill, A.T., 2013. Erosion source discrimination in a rural Australian catchment using compound-specific isotope analysis (CSIA). *Hydrological Processes* 27, 923-932.
- Hardy, F., Bariteau, L., Lorrain, S., Theriault, I., Gagnon, G., Messier, D. and Rougerie, J.F., 2010. Geochemical tracing and spatial evolution of the sediment bed load of the Romaine River, Quebec, Canada. *CATENA* 81, 66-76.
- Hatfield, R.G., 2014. Particle size-specific magnetic measurements as a tool for enhancing our understanding of the bulk magnetic properties of sediments. *Minerals* 4, 758-787.
- Hatfield, R.G. and Maher, B.A., 2008. Suspended sediment characterization and tracing using a magnetic fingerprinting technique: Bassenthwaite Lake, Cumbria, UK. *The Holocene* 18, 105-115.
- Hatfield, R.G. and Maher, B.A., 2009. Fingerprinting upland sediment sources: particle size-specific magnetic linkages between soils, lake sediments and suspended sediments. *Earth Surface Processes and Landforms* 34, 1359-1373.
- He, Q. and Owens, P.N., 1995. Determination of suspended sediment provenance using caesium-137, unsupported lead-210 and radium-226: a numerical mixing model approach. In: I. Foster, A. Gurnell and B. Webb (Editors), *Sediment and Water Quality in River Catchments*, Wiley, Chichester, 207-227.
- He, Q. and Walling, D.E., 1996. Interpreting particle size effects in the adsorption of  $^{137}\text{Cs}$  and unsupported  $^{210}\text{Pb}$  by mineral soils and sediments. *Journal of Environmental Radioactivity* 30, 117-137.
- Horowitz, A.J., 1991. A primer on sediment-trace element chemistry, 2nd edition. Open-File Report 91-76, United States Geological Survey; Books and Open-File Reports Section, Denver, Colorado, 136 pp.
- Horowitz, A.J. and Elrick, K.A., 1987. The relation of stream sediment surface area, grain size and composition to trace element chemistry. *Applied Geochemistry* 2, 437-451.
- Hughes, A.O., Olley, J.M., Croke, J.C. and McKergow, L.A., 2009. Sediment source changes over the last 250 years in a dry-tropical catchment, central Queensland, Australia. *Geomorphology* 104, 262-275.
- IAEA, 2014. Guidelines for Using Fallout Radionuclides to Assess Erosion and Effectiveness of Soil Conservation Strategies, IAEA (International Atomic Energy Agency) publication, IAEA-TECDOC-1741. Vienna, Austria, 213 pp.

934 Jagercikova, M., Cornu, S., Le Bas, C. and Evrard, O., 2015. Vertical distributions of  $^{137}\text{Cs}$  in soils: a meta-  
935 analysis. *Journal of Soils and Sediments* 15, 81-95.

936 Keil, R.G. and Mayer, L.M., 2014. 12.12 - Mineral Matrices and Organic Matter A2 - Holland, Heinrich D. In:  
937 K.K. Turekian (Editor), *Treatise on Geochemistry* (Second Edition). Elsevier, Oxford, 337-359.

938 Kersten, M. and Smedes, F., 2002. Normalization procedures for sediment contaminants in spatial and  
939 temporal trend monitoring. *Journal of Environmental Monitoring* 4, 109-115.

940 Klages, M.G. and Hsieh, Y.P., 1975. Suspended solids carried by the Gallatin River of southwestern  
941 Montana: II. Using minerology for inferring sources. *Journal of Environmental Quality* 4, 68-73.

942 Koiter, A., Lobb, D., Owens, P., Petticrew, E., Tiessen, K.D. and Li, S., 2013a. Investigating the role of  
943 connectivity and scale in assessing the sources of sediment in an agricultural watershed in the  
944 Canadian prairies using sediment source fingerprinting. *Journal of Soils and Sediments* 13, 1676-  
945 1691.

946 Koiter, A.J., Owens, P.N., Petticrew, E.L. and Lobb, D.A., 2013b. The behavioural characteristics of sediment  
947 properties and their implications for sediment fingerprinting as an approach for identifying  
948 sediment sources in river basins. *Earth-Science Reviews* 125, 24-42.

949 Koiter, A.J., Owens, P.N., Petticrew, E.L. and Lobb, D.A., 2015. The role of gravel channel beds on the  
950 particle size and organic matter selectivity of transported fine-grained sediment: implications for  
951 sediment fingerprinting and biogeochemical flux research. *Journal of Soils and Sediments* 15,  
952 2174-2188.

953 Koiter, A.J., Owens, P.N., Petticrew, E.L. and Lobb, D.A., 2017. The role of soil surface properties on the  
954 particle size and carbon selectivity of interrill erosion in agricultural landscapes. *CATENA* 153, 194-  
955 206.

956 Kraushaar, S., Schumann, T., Ollesch, G., Schubert, M., Vogel, H.-J. and Siebert, C., 2015. Sediment  
957 fingerprinting in northern Jordan: element-specific correction factors in a carbonatic setting.  
958 *Journal of Soils and Sediments* 15, 2155-2173.

959 Krumbein, W.C. and Sloss, L.L., 1951. Stratigraphy and sedimentation. *Soil Science* 71, 401.

960 Laceby, J.P., Huon, S., Onda, Y. and Evrard, O., 2016. Do forests represent a long-term source of  
961 contaminated particulate matter in the Fukushima Prefecture? *Journal of Environmental*  
962 *Management* 183, 742–753.

963 Laceby, J.P., McMahon, J., Evrard, O. and Olley, J., 2015a. A comparison of geological and statistical  
964 approaches to element selection for sediment fingerprinting. *Journal of Soils and Sediments* 15,  
965 2117-2131.

966 Laceby, J.P. and Olley, J., 2015. An examination of geochemical modelling approaches to tracing sediment  
967 sources incorporating distribution mixing and elemental correlations. *Hydrological Processes* 29,  
968 1669-1685

969 Laceby, J.P., Olley, J., Pietsch, T.J., Sheldon, F. and Bunn, S.E., 2015b. Identifying subsoil sediment sources  
970 with carbon and nitrogen stable isotope ratios. *Hydrological Processes* 29, 1956-1971.

971 Le Cloarec, M.F., Bonte, P.H., Lestel, L., Lefèvre, I. and Ayrault, S., 2011. Sedimentary record of metal  
972 contamination in the Seine River during the last century. *Physics and Chemistry of the Earth, Parts*  
973 *A/B/C* 36, 515-529.

974 Le Gall, M., Evrard, O., Foucher, A., Laceby, J.P., Salvador-Blanes, S., Thil, F., Dapoigny, A., Lefèvre, I.,  
975 Cerdan, O. and Ayrault, S., 2016. Quantifying sediment sources in a lowland agricultural catchment  
976 pond using  $^{137}\text{Cs}$  activities and radiogenic  $^{87}\text{Sr}/^{86}\text{Sr}$  ratios. *Science of The Total Environment* 566-  
977 567, 968–980.

978 Le, M.H., Cordier, S., Lucas, C. and Cerdan, O., 2015. A faster numerical scheme for a coupled system  
979 modeling soil erosion and sediment transport. *Water Resources Research* 51, 987-1005.

980 Lewis, J., 1996. Turbidity-controlled suspended sediment sampling for runoff-event load estimation. *Water*  
981 *Resources Research* 32, 2299-2310.

982 Lomenick, T.F. and Tamura, T., 1965. Naturally Occurring Fixation of Cesium-137 on Sediments of  
 983 Lacustrine Origin. *Soil Science Society of America Journal* 29, 383-387.

984 Mabit, L., Benmansour, M., Abril, J.M., Walling, D.E., Meusburger, K., Iurian, A.R., Bernard, C., Tarján, S.,  
 985 Owens, P.N. and Blake, W.H., 2014. Fallout  $^{210}\text{Pb}$  as a soil and sediment tracer in catchment  
 986 sediment budget investigations: a review. *Earth-Science Reviews* 138, 335-351.

987 Martinez-Carreras, N., Udelhoven, T., Krein, A., Gallart, F., Iffly, J.F., Ziebel, J., Hoffmann, L., Pfister, L. and  
 988 Walling, D.E., 2010. The use of sediment colour measured by diffuse reflectance spectrometry to  
 989 determine sediment sources: Application to the Attert River catchment (Luxembourg). *Journal of*  
 990 *Hydrology* 382, p 49-63.

991 Matisoff, G., Bonniwell, E.C. and Whiting, P.J., 2002. Soil erosion and sediment sources in an Ohio  
 992 watershed using Beryllium-7, Cesium-137, and Lead-210. *Journal of Environmental Quality* 31, 54-  
 993 61.

994 Matisoff, G., Wilson, C.G. and Whiting, P.J., 2005. The  $^7\text{Be}/^{210}\text{Pb}_{\text{xs}}$  ratio as an indicator of suspended  
 995 sediment age or fraction new sediment in suspension. *Earth Surface Processes and Landforms* 30,  
 996 1191-1201.

997 Mayer, L., 1999. Extent of coverage of mineral surfaces by organic matter in marine sediments. *Geochimica*  
 998 *et Cosmochimica Acta* 63, 207-215.

999 McCulloch, M., Fallon, S., Wyndham, T., Hendy, E., Lough, J. and Barnes, D., 2003. Coral record of increased  
 1000 sediment flux to the inner Great Barrier Reef since European settlement. *Nature* 421, 727-730.

1001 McLaren, P. and Bowles, D., 1985. The effects of sediment transport on grain-size distributions. *Journal of*  
 1002 *Sedimentary Research* 55, 457-470.

1003 Merkus, H.G., 2009. Particle Size Measurements: Fundamentals, Practice, Quality, 17. Springer Science &  
 1004 Business Media, 532 pp.

1005 Morgan, R., 2005. Soil Erosion and Conservation, Third Edition. Blackwell Publishing, Oxford, 303 pp.

1006 Morris, G.L. and Fan, J., 1998. Reservoir Sedimentation Handbook: Design and Management of Dams,  
 1007 Reservoirs, and Watersheds for Sustainable Use. McGraw Hill Professional, New York, 848 pp.

1008 Moss, A. and Walker, P., 1978. Particle transport by continental water flows in relation to erosion,  
 1009 deposition, soils, and human activities. *Sedimentary Geology* 20, 81-139.

1010 Motha, J.A., Wallbrink, P.J., Hairsine, P.B. and Grayson, R.B., 2002. Tracer properties of eroded sediment  
 1011 and source material. *Hydrological Processes* 16, 1983-2000.

1012 Motha, J.A., Wallbrink, P.J., Hairsine, P.B. and Grayson, R.B., 2003. Determining the sources of suspended  
 1013 sediment in a forested catchment in southeastern Australia. *Water Resources Research* 39, 1056.

1014 Mukundan, R., Radcliffe, D.E., Ritchie, J.C., Risse, L.M. and Mckinley, R.A., 2010. Sediment fingerprinting  
 1015 to determine the source of suspended sediment in a southern piedmont stream. *Journal of*  
 1016 *Environmental Quality* 39, 1328-1337.

1017 Mukundan, R., Walling, D.E., Gellis, A.C., Slattery, M.C. and Radcliffe, D.E., 2012. Sediment Source  
 1018 Fingerprinting: Transforming From a Research Tool to a Management Tool. *Journal of the*  
 1019 *American Water Resources Association* 48, 1241-1257.

1020 Murray, A., Stanton, R., Olley, J. and Morton, R., 1993a. Determining the origins and history of  
 1021 sedimentation in an underground river system using natural and fallout radionuclides. *Journal of*  
 1022 *Hydrology* 146, 341-359.

1023 Murray, A.S., Olive, L.J., Olley, J.M., Caitcheon, G.G., Wasson, R.J. and Wallbrink, P.J., 1993b. Tracing the  
 1024 Source of Suspended Sediment in the Murrumbidgee River, Australia. In: N.E. Peters, E. Hoehn, C.  
 1025 Leibundgut, N. Tase and D.E. Walling (Editors), *Tracers in Hydrology*, IAHS Publication No. 215,  
 1026 IAHS Press, Wallingford, 293-302.

1027 Natelhoffer, K.J. and Fry, B., 1988. Controls on natural nitrogen-15 and carbon-13 abundances in forest  
 1028 soil organic matter. *Soil Science Society of America Journal* 52, 1633-1640.

1029 Navratil, O., Evrard, O., Esteves, M., Legout, C., Ayrault, S., Némery, J., Mate-Marin, A., Ahmadi, M.,  
1030 Lefèvre, I. and Poirel, A., 2012. Temporal variability of suspended sediment sources in an alpine  
1031 catchment combining river/rainfall monitoring and sediment fingerprinting. *Earth Surface*  
1032 *Processes and Landforms* 37, 828-846.

1033 Oldfield, F., 1991. Environmental magnetism — A personal perspective. *Quaternary Science Reviews* 10,  
1034 73-85.

1035 Oldfield, F., Maher, B., Donoghue, J. and Pierce, J., 1985. Particle-size related, mineral magnetic source  
1036 sediment linkages in the Rhode River catchment, Maryland, USA. *Journal of the Geological Society*  
1037 142, 1035-1046.

1038 Olley, J., Burton, J., Smolders, K., Pantus, F. and Pietsch, T., 2013. The application of fallout radionuclides  
1039 to determine the dominant erosion process in water supply catchments of subtropical South-east  
1040 Queensland, Australia. *Hydrological Processes* 27, 885-895.

1041 Olley, J. and Caitcheon, G., 2000. Major element chemistry of sediments from the Darling-Barwon river  
1042 and its tributaries: implications for sediment and phosphorus sources. *Hydrological Processes* 14,  
1043 1159-1175.

1044 Olley, J.M., 2002. Organic carbon supply to a large lowland river and implications for aquatic ecosystems.  
1045 In: F.J. Dyer, M.C. Thoms and J.M. Olley (Editors), *The Structure Function and Management*  
1046 *Implications of Fluvial Sedimentary Systems*, IAHS Publication No. 276, IAHS Press, Wallingford,  
1047 27-33.

1048 Olley, J.M., Caitcheon, G.G., Hancock, G. and Wallbrink, P.J., 2001. *Tracing and Dating Techniques for*  
1049 *Sediment and Associated Substances*. CSIRO Land and Water, Canberra.

1050 Olley, J.M. and Murray, A.S., 1994. Origins of variability in the  $^{230}\text{Th}/^{232}\text{Th}$  ratio in sediments. In: D.E. Walling  
1051 and Q. He (Editors), *Variability in Stream Erosion and Sediment Transport*, IAHS Publication No.  
1052 224, IAHS Press, Wallingford, 65-70.

1053 Ortiz, J.D., Polyak, L., Grebmeier, J.M., Darby, D., Eberl, D.D., Naidu, S. and Nof, D., 2009. Provenance of  
1054 Holocene sediment on the Chukchi-Alaskan margin based on combined diffuse spectral  
1055 reflectance and quantitative X-Ray Diffraction analysis. *Global and Planetary Change* 68, 73-84.

1056 Owens, P.N., Batalla, R.J., Collins, A.J., Gomez, B., Hicks, D.M., Horowitz, A.J., Kondolf, G.M., Marden, M.,  
1057 Page, M.J., Peacock, D.H., Petticrew, E.L., Salomons, W. and Trustrum, N.A., 2005. Fine-grained  
1058 sediment in river systems: environmental significance and management issues. *River Research and*  
1059 *Applications* 21, 693-717.

1060 Owens, P.N., Blake, W.H., Gaspar, L., Gateuille, D., Koiter, A.J., Lobb, D.A., Petticrew, E.L., Raiffarth, D.,  
1061 Smith, H.G. and Woodward, J.C., 2016. Fingerprinting and tracing the sources of soils and  
1062 sediments: earth and ocean science, geoarchaeological, forensic, and human health applications.  
1063 *Earth-Science Reviews* 162, 1-23.

1064 Owens, P.N., Blake, W.H., Giles, T.R. and Williams, N.D., 2012. Determining the effects of wildfire on  
1065 sediment sources using  $^{137}\text{Cs}$  and unsupported  $^{210}\text{Pb}$ : the role of landscape disturbances and  
1066 driving forces. *Journal of Soils and Sediments* 12, 982-994.

1067 Owens, P.N., Blake, W.H. and Petticrew, E.L., 2006. Changes in sediment sources following wildfire in  
1068 mountainous terrain: A paired-catchment approach, British Columbia, Canada. In: B. Kronvang, J.  
1069 Faganeli and N. Ogrinc (Editors), *The Interactions Between Sediments and Water*. Springer  
1070 Netherlands, 273-281.

1071 Owens, P.N., Walling, D.E. and Leeks, G.J.L., 2000. Tracing fluvial suspended sediment sources in the  
1072 catchment of the River Tweed, Scotland, using composite fingerprints and a numerical mixing  
1073 model. In: I. Foster (Editor), *Tracers in Geomorphology*. Wiley, Chichester, 291-308.

1074 Palazón, L., Gaspar, L., Latorre, B., Blake, W. and Navas, A., 2015a. Identifying sediment sources by applying  
1075 a fingerprinting mixing model in a Pyrenean drainage catchment. *Journal of Soils and Sediments*  
1076 15, 2067-2085.

1077 Palazón, L., Latorre, B., Gaspar, L., Blake, W.H., Smith, H.G. and Navas, A., 2015b. Comparing catchment  
1078 sediment fingerprinting procedures using an auto-evaluation approach with virtual sample  
1079 mixtures. *Science of The Total Environment* 532, 456-466.

1080 Panuska, J.C., Karthikeyan, K.G. and Miller, P.S., 2008. Impact of surface roughness and crusting on particle  
1081 size distribution of edge-of-field sediments. *Geoderma* 145, 315-324.

1082 Papanicolaou, A.N., Fox, J.F. and Marshall, J., 2003. Soil fingerprinting in the Palouse Basin, USA using  
1083 stable carbon and nitrogen isotopes. *International Journal of Sediment Research* 18, 278-284.

1084 Peart, M.R. and Walling, D.E. (Editors), 1986. Fingerprinting sediment source: The example of a drainage  
1085 basin in Devon, UK. *Drainage Basin Sediment Delivery*, IAHS Publication No. 159, IAHS Press,  
1086 Wallingford, 41-55 pp.

1087 Phillips, J. and Walling, D., 1995. An assessment of the effects of sample collection, storage and  
1088 resuspension on the representativeness of measurements of the effective particle size distribution  
1089 of fluvial suspended sediment. *Water Research* 29, 2498-2508.

1090 Poesen, J., 1992. Mechanisms of overland-flow generation and sediment production on loamy and sandy  
1091 soils with and without rock fragments. In: A.J. Parsons and A.D. Abrahams (Editors), *Overland Flow*  
1092 *Hydraulics and Erosion Mechanics*. Routledge, London, 275-305.

1093 Poleto, C., Merten, G.H. and Minella, J.P., 2009. The identification of sediment sources in a small urban  
1094 watershed in southern Brazil: An application of sediment fingerprinting. *Environmental*  
1095 *Technology* 30, 1145-1153.

1096 Poulenard, J., Legout, C., Némery, J., Bramorski, J., Navratil, O., Douchin, A., Fanget, B., Perrette, Y., Evrard,  
1097 O. and Esteves, M., 2012. Tracing sediment sources during floods using Diffuse Reflectance  
1098 Infrared Fourier Transform Spectrometry (DRIFTS): A case study in a highly erosive mountainous  
1099 catchment (Southern French Alps). *Journal of Hydrology* 414-415, 452-462.

1100 Poulenard, J., Perrette, Y., Fanget, B., Quetin, P., Trevisan, D. and Dorioz, J.M., 2009. Infrared spectroscopy  
1101 tracing of sediment sources in a small rural watershed (French Alps). *Science of The Total*  
1102 *Environment* 407, 2808-2819.

1103 Pulley, S., Foster, I. and Antunes, P., 2015a. The application of sediment fingerprinting to floodplain and  
1104 lake sediment cores: assumptions and uncertainties evaluated through case studies in the Nene  
1105 Basin, UK. *Journal of Soils and Sediments*, 1-23.

1106 Pulley, S., Rowntree, K. and Foster, I., 2015b. Conservatism of mineral magnetic signatures in farm dam  
1107 sediments in the South African Karoo: the potential effects of particle size and post-depositional  
1108 diagenesis. *Journal of Soils and Sediments* 15, 2387-2397.

1109 Rawlins, B.G., Turner, G., Mounteney, I. and Wildman, G., 2010. Estimating specific surface area of fine  
1110 stream bed sediments from geochemistry. *Applied Geochemistry* 25, 1291-1300.

1111 Reiffarth, D.G., Petticrew, E.L., Owens, P.N. and Lobb, D.A., 2016. Sources of variability in fatty acid (FA)  
1112 biomarkers in the application of compound-specific stable isotopes (CSSIs) to soil and sediment  
1113 fingerprinting and tracing: A review. *Science of The Total Environment* 565, 8-27.

1114 Russell, M.A., Walling, D.E. and Hodgkinson, R.A., 2001. Suspended sediment sources in two small lowland  
1115 agricultural catchments in the UK. *Journal of Hydrology* 252, 1-24.

1116 Schimel, D.S., 1993. *Theory and Application of Tracers. Isotopic Techniques in Plant, Soil and Aquatic*  
1117 *Biology*. Academic Press, Inc., San Diego., 119 pp.

1118 Sherriff, S.C., Franks, S.W., Rowan, J.S., Fenton, O. and Ó'hUallacháin, D., 2015. Uncertainty-based  
1119 assessment of tracer selection, tracer non-conservativeness and multiple solutions in sediment  
1120 fingerprinting using synthetic and field data. *Journal of Soils and Sediments* 10, 2101-2116.

1121 Singh, M., Sharma, M. and Tobschall, H.J., 2005. Weathering of the Ganga alluvial plain, northern India:  
1122 implications from fluvial geochemistry of the Gomati River. *Applied Geochemistry* 20, 1-21.

1123 Singleton, A.A., Schmidt, A.H., Bierman, P.R., Rood, D.H., Neilson, T.B., Greene, E.S., Bower, J.A. and  
1124 Perdrial, N., 2017. Effects of grain size, mineralogy, and acid-extractable grain coatings on the

1125 distribution of the fallout radionuclides <sup>7</sup>Be, <sup>10</sup>Be, <sup>137</sup>Cs, and <sup>210</sup>Pb in river sediment.  
1126 *Geochimica et Cosmochimica Acta* 197, 71-86.

1127 Slattery, M.C., Burt, T.P. and Walden, J., 1995. The application of mineral magnetic measurements to  
1128 quantify within-storm variations in suspended sediment sources. In: C. Leibundgut (Editor), *Tracer*  
1129 *Technologies for Hydrological Systems*, IAHS Publication No. 229, IAHS Press, Wallingford, 143-  
1130 152.

1131 Smith, H.G. and Blake, W.H., 2014. Sediment fingerprinting in agricultural catchments: A critical re-  
1132 examination of source discrimination and data corrections. *Geomorphology* 204, 177-191.

1133 Smith, H.G., Blake, W.H. and Owens, P.N., 2013. Discriminating fine sediment sources and the application  
1134 of sediment tracers in burned catchments: a review. *Hydrological Processes* 27, 943-958.

1135 Smith, H.G., Blake, W.H. and Taylor, A., 2014. Modelling particle residence times in agricultural river basins  
1136 using a sediment budget model and fallout radionuclide tracers. *Earth Surface Processes and*  
1137 *Landforms* 39, 1944-1959.

1138 Smith, H.G., Sheridan, G.J., Lane, P.N., Noske, P.J. and Heijnis, H., 2011. Changes to sediment sources  
1139 following wildfire in a forested upland catchment, southeastern Australia. *Hydrological Processes*  
1140 25, 2878-2889.

1141 Smith, J., 1999. An introduction to the magnetic properties of natural materials. *Environmental*  
1142 *Magnetism: A Practical Guide*. Technical Guide 6, 5-34.

1143 Stone, M., Collins, A., Silins, U., Emelko, M. and Zhang, Y., 2014. The use of composite fingerprints to  
1144 quantify sediment sources in a wildfire impacted landscape, Alberta, Canada. *Science of The Total*  
1145 *Environment* 473, 642-650.

1146 Stone, P. and Walling, D., 1997. Particle size selectivity considerations in suspended sediment budget  
1147 investigations. In: R.D. Evans, J. Wisniewski and J.R. Wisniewski (Editors), *The Interactions Between*  
1148 *Sediments and Water*, Springer, Netherlands, 63-70.

1149 Stout, J.C., Belmont, P., Schottler, S.P. and Willenbring, J.K., 2014. Identifying sediment sources and sinks  
1150 in the Root River, Southeastern Minnesota. *Annals of the Association of American Geographers*  
1151 104, 20-39.

1152 Stumm, W. and Morgan, J.J., 2012. *Aquatic Chemistry: Chemical Equilibria and rates in Natural Waters*.  
1153 John Wiley & Sons, Chichester, 1040 pp.

1154 Szava-Kovats, R.C., 2008. Grain-size normalization as a tool to assess contamination in marine sediments:  
1155 Is the < 63µm fraction fine enough? *Marine Pollution Bulletin* 56, 629-632.

1156 Tamura, T., 1964. Consequences of activity release: selective sorption reactions of cesium with soil  
1157 minerals. *Nuclear Safety* 5, 262-268.

1158 Taylor, A., Blake, W.H. and Keith-Roach, M.J., 2014. Estimating Be-7 association with soil particle size  
1159 fractions for erosion and deposition modelling. *Journal of Soils and Sediments* 14, 1886-1893.

1160 Taylor, A., Blake, W.H., Smith, H.G., Mabit, L. and Keith-Roach, M.J., 2013. Assumptions and challenges in  
1161 the use of fallout beryllium-7 as a soil and sediment tracer in river basins. *Earth-Science Reviews*  
1162 126, 85-95.

1163 Tiecher, T., Caner, L., Minella, J.P.G., Bender, M.A. and dos Santos, D.R., 2016. Tracing sediment sources in  
1164 a subtropical rural catchment of southern Brazil by using geochemical tracers and near-infrared  
1165 spectroscopy. *Soil and Tillage Research* 155, 478-491.

1166 Vale, S.S., Fuller, I.C., Procter, J.N., Basher, L.R. and Smith, I.E., 2016. Characterization and quantification  
1167 of suspended sediment sources to the Manawatu River, New Zealand. *Science of The Total*  
1168 *Environment* 543, 171-186.

1169 Vercruysse, K., Grabowski, R.C. and Rickson, R.J., 2017. Suspended sediment transport dynamics in rivers:  
1170 Multi-scale drivers of temporal variation. *Earth-Science Reviews* 166, 38-52.

1171 Viparelli, E., Wesley Lauer, J., Belmont, P. and Parker, G., 2013. A numerical model to develop long-term  
1172 sediment budgets using isotopic sediment fingerprints. *Computers & Geosciences* 53, 114-122.



1173 Walden, J. and Slattery, M.C., 1993. Verification of a simple gravity technique for separation of particle  
1174 size fractions suitable for mineral magnetic analysis. *Earth Surface Processes and Landforms* 18,  
1175 829-833.

1176 Wall, G.J. and Wilding, L.P., 1975. Minerology and Related Parameters of Fluvial Suspended Sediments in  
1177 Northwestern Ohio. *Journal of Environmental Quality* 5, 168-173.

1178 Wallbrink, P. and Murray, A., 1996a. Distribution and variability of <sup>7</sup>Be in soils under different surface  
1179 cover conditions and its potential for describing soil redistribution processes. *Water Resources*  
1180 *Research* 32, 467-476.

1181 Wallbrink, P.J. and Murray, A.S., 1993. Use of fallout radionuclides as indicators of erosion processes.  
1182 *Hydrological Processes* 7, 297-304.

1183 Wallbrink, P.J. and Murray, A.S., 1996b. Determining Soil Loss Using The Inventory Ratio Of Excess-Lead  
1184 210 To Cesium-137. *Soil Science Society of America Journal* 60, 1201-1208.

1185 Wallbrink, P.J., Murray, A.S., Olley, J.M. and Olive, L.J., 1998. Determining sources and transit times of  
1186 suspended sediment in the Murrumbidgee River, New South Wales, Australia, using fallout <sup>137</sup>Cs  
1187 and <sup>210</sup>Pb. *Water Resources Research* 34, 879-887.

1188 Wallbrink, P.J., Olley, J.M. and Murray, A.S., 1999. Relating suspended sediment to its original soil depth  
1189 using fallout radionuclides. *Soil Science Society of America Journal* 63, 369-378.

1190 Walling, D. and Moorehead, P., 1989. The particle size characteristics of fluvial suspended sediment: an  
1191 overview. *Hydrobiologia* 176, 125-149.

1192 Walling, D., Peart, M., Oldfield, F. and Thompson, R., 1979. Suspended sediment sources identified by  
1193 magnetic measurements. *Nature* 281, 110-113.

1194 Walling, D.E., 2003. Using environmental radionuclides as tracers in sediment budget investigations. In:  
1195 D.E. Walling and T. Fergus (Editors), *Erosion and Sediment Transport Measurement in Rivers:*  
1196 *Technological and Methodological Advances*, IAHS Publication No. 283, IAHS Press, Wallingford,  
1197 57-78.

1198 Walling, D.E., 2005. Tracing suspended sediment sources in catchments and river systems. *Science of The*  
1199 *Total Environment* 344, 159-184.

1200 Walling, D.E. and Collins, A.L., 2016. Fine sediment transport and management. In: D.J. Gilvear, M.T.  
1201 Greenwood, M.C. Thoms and P.J. Wood (Editors), *River Science: Research and Management for*  
1202 *the 21st Century*, John Wiley & Sons, Chichester, 37-60.

1203 Walling, D.E. and Kane, P., 1984. Suspended sediment properties and their geomorphological significance.  
1204 In: T.P. Burt and D.E. Walling (Editors), *Catchment Experiments in Fluvial Geomorphology*, Geo  
1205 Books, Norwich, 311-334.

1206 Walling, D.E., Owens, P.N. and Leeks, G.J.L., 1999. Fingerprinting suspended sediment sources in the  
1207 catchment of the River Ouse, Yorkshire, UK. *Hydrological Processes* 13, 955-975.

1208 Walling, D.E., Owens, P.N., Waterfall, B.D., Leeks, G.J. and Wass, P.D., 2000. The particle size characteristics  
1209 of fluvial suspended sediment in the Humber and Tweed catchments, UK. *Science of The Total*  
1210 *Environment* 251, 205-222.

1211 Walling, D.E. and Woodward, J.C., 1992. Use of radiometric fingerprints to derive information on  
1212 suspended sediment sources In: J. Bogen, D.E. Walling and T. Day (Editors), *Erosion and Sediment*  
1213 *Transport Monitoring Programmes in River Basins*, IAHS Publication No. 210, IAHS Press,  
1214 Wallingford, 153-164.

1215 Walling, D.E., Woodward, J.C. and Nicholas, A.P., 1993. A multi-parameter approach to fingerprinting  
1216 suspended-sediment sources. In: N.E. Peters, E. Hoehn, C. Leibundgut, N. Tase and D.E. Walling  
1217 (Editors), *Tracers in Hydrology*, IAHS Publication No. 215, IAHS Press, Wallingford, 329-338.

1218 Wischmeier, W.H. and Smith, D.D., 1978. Predicting Rainfall Erosion Losses-A guide to Conservation  
1219 Planning. *Agricultural Handbook Number 537*, United States Department of Agriculture, Science  
1220 and Education Administration, Hyattsville, Maryland, 58 pp.

1221 Wohl, E., Bledsoe, B.P., Jacobson, R.B., Poff, N.L., Rathburn, S.L., Walters, D.M. and Wilcox, A.C., 2015. The  
1222 Natural Sediment Regime in Rivers: Broadening the Foundation for Ecosystem Management.  
1223 BioScience 65, 358-371.

1224 Wynn, J.G., Bird, M.I. and Wong, V.N.L., 2005. Rayleigh distillation and the depth profile of  $^{13}\text{C}/^{12}\text{C}$  ratios  
1225 of soil organic carbon from soils of disparate texture in Iron Range National Park, Far North  
1226 Queensland, Australia. *Geochimica et Cosmochimica Acta* 69, 1961-1973.

1227 Yamashiki, Y., Onda, Y., Smith, H.G., Blake, W.H., Wakahara, T., Igarashi, Y., Matsuura, Y. and Yoshimura,  
1228 K., 2014. Initial flux of sediment-associated radiocesium to the ocean from the largest river  
1229 impacted by Fukushima Daiichi Nuclear Power Plant. *Scientific Reports* 4.

1230 Zhang, X.J. and Liu, B., 2016. Using multiple composite fingerprints to quantify fine sediment source  
1231 contributions: A new direction. *Geoderma* 268, 108-118.

1232